

Phase coherence and superfluid-insulator transition in a disordered Bose-Einstein condensate

Yong P. Chen,^{1,2,*} J. Hitchcock,¹ D. Dries,¹ M. Junker,¹ C. Welford,¹ and R. G. Hulet¹

¹Department of Physics and Astronomy and Rice Quantum Institute, Rice University, 6100 Main Street, Houston, Texas 77005, USA

²The Richard E. Smalley Institute for Nanoscale Science and Technology, Rice University, 6100 Main Street, Houston, Texas 77005, USA

(Received 26 October 2007; published 31 March 2008)

We have studied the effects of a disordered optical potential on the transport and phase coherence of a Bose-Einstein condensate (BEC) of ⁷Li atoms. At moderate disorder strengths (V_D), we observe inhibited transport and damping of dipole excitations, while in time-of-flight images, random but reproducible interference patterns are observed. *In situ* images reveal that the appearance of interference is correlated with density modulation, without complete fragmentation. At higher V_D , the interference contrast diminishes as the BEC fragments into multiple pieces with little phase coherence.

DOI: [10.1103/PhysRevA.77.033632](https://doi.org/10.1103/PhysRevA.77.033632)

PACS number(s): 03.75.Hh, 03.75.Kk, 64.60.Cn

The behavior of a superfluid or a superconductor in the presence of disorder is of fundamental interest. A superfluid can flow without friction around obstacles, and a superconductor can have zero resistance despite material defects. On the other hand, disorder is able to localize particles, resulting in an insulating state [1]. Experimentally, disorder-induced superfluid/superconductor to insulator transitions (SIT) have been probed in many systems, including superfluid helium in porous media [2], thin-film and granular superconductors [3,4], and random Josephson junction arrays [5]. While many believe that such a SIT is a quantum phase transition driven by quantum fluctuations, it remains a central task to understand exactly how the superfluid/superconducting order parameter, which consists both an amplitude and a phase, may be destroyed with increasing disorder. Numerous fundamental questions remain, such as the nature of the insulator, the fate of phase coherence throughout the transition, and the possibility of intermediate metallic phases [3,6,7].

Cold atoms, with their intrinsic cleanliness coupled with remarkable controllability of physical parameters, have emerged as exceptional systems to study various condensed matter problems. Recently, several experiments [8–11] have studied ⁸⁷Rb condensates in *random* optical potentials and observed, for example, damping of collective excitations [8] and inhibition of expansion [9–11] due to disorder. Another experiment [12] has examined a Bose-Einstein condensate (BEC) in an incommensurate (*quasirandom*) optical lattice in order to investigate a possible “Bose-glass” phase [13]. Experiments with disordered atomic quantum gases may provide unique insights into disordered quantum systems and may uncover a rich variety of quantum phases [14].

Here we report experiments on a BEC of interacting ⁷Li atoms subject to a well-controlled disordered potential. While we corroborate previous transport measurements [8–11], we have also probed the ground-state density distribution and phase coherence of the disordered BEC by performing both *in situ* and time-of-flight (TOF) imaging. While disorder inhibits transport of the BEC, *reproducible* TOF interference patterns are observed for intermediate disorder

strengths V_D , reflecting an underlying phase coherence in the disordered BEC. At stronger V_D , the interference contrast diminishes as the BEC fragments into a “granular” condensate, which is expected to have no phase coherence.

We create an optically trapped BEC of ⁷Li atoms in the $m_f = -1/2$, $m_f = 3/2$ state at a bias magnetic field $B \approx 720$ G using procedures essentially similar to those described in [15]. The bias field tunes the scattering length a_s via a Feshbach resonance to a large positive value ($\sim 200a_0$, where a_0 is the Bohr radius), enabling efficient evaporative cooling [15,16]. The optical dipole trap is produced by focusing a 1030 nm Yb:YAG laser beam to a Gaussian $1/e^2$ intensity radius of 34 μm . In the following, we label the axial direction (along which the optical trap beam propagates) as z and the two radial directions as x and y . At the final trap depth, the radial trap frequencies are $\omega_x = \omega_y \approx 2\pi \times 180$ Hz and the axial trap frequency is $\omega_z \approx 2\pi \times 3.6$ Hz. Part of the axial confinement is contributed by a residual curvature from the bias magnetic field. At this trap depth, the BEC contains $\sim 5 \times 10^5$ atoms with no discernible thermal component. The corresponding chemical potential $\mu \approx h \times 1$ kHz, where h is Planck’s constant.

To create the disordered potential, we pass another beam, derived from the same laser as used for the optical trap, through a diffusive plate. This results in laser speckle [17] similar to that used in [8,10]. This beam propagates along the x direction. Because of the optical dipole force, the speckle’s spatially varying intensity pattern gives rise to a disordered potential $V(x, y, z)$ for the atoms. Figure 1(a) shows a two-dimensional (2D) cross-section $V(0, y, z)$ as imaged by a camera, while the main panel shows an axial cut $V(z) = V(0, 0, z)$ of the disordered potential through the center ($x = y = 0$) of the atomic cloud. The disorder strength V_D is proportional to the intensity of the speckle laser and is defined as the standard deviation of the mean of $V(z)$. The disorder correlation length along the z direction, σ_z , is defined by fitting the autocorrelation function $\langle V(z)V(z+\Delta z) \rangle$ to $e^{-2\Delta z^2/\sigma_z^2}$ [Fig. 1(b)]; we find $\sigma_z \approx 15$ μm , which is much greater than the condensate healing length of ~ 1 μm . Since $\sigma_x \gg \sigma_y = \sigma_z$ [10,17] and the radial Thomas-Fermi radius of the BEC is less than 10 μm , to a good approximation the BEC is subject to a 1D disordered potential $V(z) = V(0, 0, z)$ along its axial direction.

*Current address: Purdue University, 525 Northwestern Ave., West Lafayette, IN 47907, USA.




FIG. 1. (Color) Typical 1D disordered potential $V(z)$. (a) False-color surface plot of the disordered speckle potential $V(y, z)$. The pixel size corresponds to $2.2 \mu\text{m}$ in the plane of the atoms. (b) Autocorrelation function of $V(z)$. The line is a fit to $e^{-2\Delta z^2/\sigma^2}$, within a scaling factor. As expected for fully developed speckle, the intensity follows a negative exponential probability distribution.

After producing a pure BEC (to within our resolution), the disordered potential is ramped on in ~ 100 ms [18] to a specific value of V_D , which we express in terms of μ . The BEC is then held in the disordered and trap potentials while various experiments are performed, as described below. The final state is probed by absorption imaging (also along the x direction) of the atomic cloud released from the disordered and optical trap potentials or by *in situ* phase-contrast imaging [19].

We first describe data characterizing the transport properties of the disordered BEC. In one type of measurement, we *slowly* ramp on a magnetic field gradient along the z direction, which offsets the center of the harmonic trap from $z=0$ to $z=-d$. The disordered potential is kept stationary during the trap offset. As shown in Figs. 2(a) and 2(b), without disorder the center of the atomic cloud follows the trap. For intermediate disorder strength the cloud lags behind the new trap center and is stretched, as shown in Fig. 2(c). For stronger disorder, as shown in Fig. 2(d), the cloud is pinned at its initial position and does not respond to the offset. We plot the cloud center as a function of V_D for four different offset distances d in Fig. 2. The cloud is pinned and does not respond to trap offset for $V_D \geq 0.8\mu$, within our experimental time scale and uncertainty.

We have also studied the effect of disorder on the dipole oscillation of the trapped BEC (Fig. 3). In contrast to the slow offset experiment discussed above, here the harmonic trap center is *abruptly* (2 ms) offset and the cloud evolves in the presence of the shifted trap and stationary disorder for a variable time before being released for imaging. Without disorder, the condensate undergoes undamped dipole oscillations in the trap with frequency ω_z . Even disorder as small as $V_D \sim 0.1 \mu$, however, damps these oscillations. Increasing the disorder strength to $V_D \geq 0.4 \mu$ causes the dipole motion to be overdamped, until when $V_D \geq \mu$, the cloud becomes pinned at its initial position to within our experimental resolution.




FIG. 2. (Color) Effect of disorder on slow transport of the BEC. The trap center is offset to the left for a distance d over a time interval of 700 ms. (a) Initial state of the BEC, without disorder or offset. The center of this initial cloud defines the reference position “0.” (b)–(d) Result of offsetting the trap by $d \approx 1400 \mu\text{m}$, with the cloud subjected to a disordered potential of various V_D : (b) $V_D=0$, (c) $V_D \approx 0.3 \mu$, and (d) $V_D \approx \mu$. Images (a)–(d) are obtained by absorption imaging taken with a 1-ms TOF after the cloud is abruptly released from the disordered and optical trap potentials. Main panel shows the center by weight of the cloud as a function of V_D , for several offsets d . V_D and μ have uncertainties of $\sim 20\%$.

In addition to transport measurements, we have performed *in situ* [20] and TOF imaging on the disordered BEC, yielding information on both its density and phase. Figures 4(a) and 4(b) show *in situ* and corresponding TOF images taken 8 ms after releasing the cloud. For intermediate V_D [Figs. 4(b) and 4(h)] striking random fringes, which we interpret as matter wave interference, develop after sufficiently long TOF expansion of the BEC following release from the optical potentials. We note that at such V_D , the corresponding *in situ*




FIG. 3. (Color) Effect of disorder on dipole excitation of the condensate. Data show the cloud center as a function of time following a sudden axial offset of the trap for various disorder strengths V_D (shown in two separate panels for clarity). The lines are fits to damped dipole oscillations.




FIG. 4. (Color) *In situ* profiles and matter wave interference. (a) *In situ* image for $V_D \approx 0.5 \mu$ and (b) TOF image for $V_D \approx 0.5 \mu$. The TOF 2D image uses a different color scale from the one used for the *in situ* 2D image, and the displayed aspect ratio of the *in situ* images is reduced by 50% in order to enhance the transverse size for visibility. (c)–(j) *In situ* column-density profiles on left and corresponding TOF profiles on right: (c),(d) $V_D = 0$; (e),(f) $V_D \approx 0.3 \mu$; (g),(h) $V_D \approx 0.5 \mu$; and (i),(j) $V_D \approx 1.0 \mu$. Solid red lines are fits to Thomas-Fermi distributions. $\omega_z \approx 2\pi \times 2.8$ Hz for data shown in Figs. 4 and 5.

images [Figs. 4(a) and 4(g)] are consistent with the cloud still being *connected*. By increasing V_D above μ , the disordered BEC becomes *fragmented* as shown in the *in situ* profile, Fig. 4(i), and the fringe contrast observed in TOF diminishes, as shown in Fig. 4(j). To quantify the dependence of the TOF fringe pattern on V_D , we plot both the TOF and *in situ* contrast versus V_D/μ in Fig. 5. The TOF fringe contrast peaks at $V_D \approx 0.5 \mu$, while the contrast in the *in situ* images increases up to the highest V_D investigated.




FIG. 5. (Color online) Contrast of the matter wave interference pattern in 8-ms TOF images (solid diamonds) and contrast of *in situ* density fluctuations (open circles) as a function of V_D/μ . The contrast is extracted from the central axial cut of the column density as its mean deviation from the Thomas-Fermi fit normalized by the peak value of the fit.

Similar, but less well resolved fringes in TOF images have been reported previously [8,11]. Our highly elongated BEC facilitates a systematic and quantitative study of the fringes. We find that the positions of these irregularly spaced fringes are *reproducible* in repeated measurements, even with different holding times of the BEC in the disordered potential. During preparation of this paper, we became aware of a closely related paper [21], which also reports reproducible fringes in TOF images of a disordered BEC. The reproducibility suggests that these fringes are unlikely due to some initial (before release) phase fluctuations [22] in the disordered BEC or due to interference of a *few* BECs separated by high potential barriers with no well-defined relative phase between them, as in either of these scenarios the fringe positions are not expected to be reproducible.

Numerical simulations based on the Gross-Pitaevskii equation [11,21], have shown that such interference can occur when moderate disorder-induced density fluctuations cause different parts of a *phase-coherent* BEC to expand with different velocities and overlap. As V_D is increased from 0, the contrast of the interference is expected to increase, consistent with the data shown in Fig. 5 at small V_D . When V_D becomes sufficiently large, however, the BEC fragments into multiple pieces, as is seen in the *in situ* image, Fig. 4(i). An array of *randomly spaced* condensates should not produce visible interference in TOF [23,24], again consistent with our observation of diminishing fringe contrast at high V_D [Fig. 4(j)].

At high V_D , we observe completely inhibited transport, indicating a transition to an insulator. This was seen in previous transport experiments, where the nature of the insulator is inferred to be that of a fragmented BEC [8–10]. Our *in situ* images give the first direct observation of fragmentation of a highly disordered BEC. Because of the exponentially suppressed Josephson tunneling between fragments due to high potential barriers, the fragmented BEC is expected to have large phase fluctuations and no phase coherence [25]. This regime is analogous to a granular superconductor [3,4], which is an insulator without global phase coherence.

Dipole oscillations are damped at moderate V_D , as shown in Fig. 3, and in agreement with previous observations [8]. Excitations such as solitons, vortices, and phase slips have been suggested [27] as possible mechanisms giving rise to damping. Our *in situ* as well as TOF data at moderate V_D suggest a disordered BEC that is still connected (not yet fragmented) and phase coherent in order to give a reproducible interference pattern. Such a state may provide insight into the physics of “homogenous” disordered superconducting films, where there has been tremendous interest regarding the form of the order parameter, possibilities of metallic states, and the fate of phase coherence and phase stiffness [28] during the SIT [3,7]. Damped dipole oscillations were also recently studied for a BEC in a periodic potential [29] in the context of a possible “Bose metal” phase [7].

Strong repulsive interactions are believed to preclude Anderson localization (AL) [11,25] in a BEC such as ours and those in previous experiments where the condensate healing length is small in comparison to σ_z [8–11]. Interactions, however, can be significantly reduced in ^7Li due to

an extremely gradual zero crossing in the B dependence of a_s [15,16], creating more favorable conditions to observe AL. Such experiments are currently underway. A similar zero crossing has been identified in ^{39}K [26].

Financial support of this work was provided by NSF, ONR, the Welch Foundation (Grant No. C-1133), and the Keck Foundation. We thank A. Aspect, L. Sanchez-Palencia, and T. A. Corcovilos for helpful discussions.

-
- [1] P. W. Anderson, Phys. Rev. **109**, 1492 (1958); P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985).
- [2] J. D. Reppy, J. Low Temp. Phys. **87**, 205 (1992).
- [3] A. M. Goldman and N. Markovic, Phys. Today **51**(11), 39 (1998); Y. Dubi, Y. Meir, and Y. Avishai, Nature (London) **449**, 876 (2007).
- [4] I. S. Beloborodov, A. V. Lopatin, V. M. Vinokur, and K. B. Efetov, Rev. Mod. Phys. **79**, 469 (2007).
- [5] E. Granato and J. M. Kosterlitz, Phys. Rev. B **33**, 6533 (1986); M. Y. Choi, J. S. Chung, and D. Stroud, *ibid.* **35**, 1669 (1987).
- [6] M. P. A. Fisher, Phys. Rev. Lett. **65**, 923 (1990).
- [7] P. Phillips and D. Dalidovich, Science **302**, 243 (2003).
- [8] J. E. Lye, L. Fallani, M. Modugno, D. S. Wiersma, C. Fort, and M. Inguscio, Phys. Rev. Lett. **95**, 070401 (2005).
- [9] C. Fort, L. Fallani, V. Guarrera, J. E. Lye, M. Modugno, D. S. Wiersma, and M. Inguscio, Phys. Rev. Lett. **95**, 170410 (2005).
- [10] D. Clément, A. F. Varon, M. Hugbart, J. A. Retter, P. Bouyer, L. Sanchez-Palencia, D. M. Gangardt, G. V. Shlyapnikov, and A. Aspect, Phys. Rev. Lett. **95**, 170409 (2005); D. Clément *et al.*, New J. Phys. **8**, 165 (2006).
- [11] T. Schulte, S. Drenkelforth, J. Kruse, W. Ertmer, J. Arlt, K. Sacha, J. Zakrzewski, and M. Lewenstein, Phys. Rev. Lett. **95**, 170411 (2005).
- [12] L. Fallani, J. E. Lye, V. Guarrera, C. Fort, and M. Inguscio, Phys. Rev. Lett. **98**, 130404 (2007); J. E. Lye, L. Fallani, C. Fort, V. Guarrera, M. Modugno, D. S. Wiersma, and M. Inguscio, Phys. Rev. A **75**, 061603(R) (2007).
- [13] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, Phys. Rev. B **40**, 546 (1989); W. Krauth, N. Trivedi, and D. Ceperley, Phys. Rev. Lett. **67**, 2307 (1991).
- [14] B. Damski, J. Zakrzewski, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **91**, 080403 (2003); A. Sanpera, A. Kantian, L. Sanchez-Palencia, J. Zakrzewski, and M. Lewenstein, *ibid.* **93**, 040401 (2004).
- [15] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, Nature (London) **417**, 150 (2002).
- [16] L. Khaykovich *et al.*, Science **296**, 1290 (2002).
- [17] J. W. Goodman, *Speckle Phenomena in Optics* (Roberts, Greenwood Village, CO, 2007).
- [18] We have checked that ramping on of the disordered potential does not produce significant heating by ramping the disordered potential back to zero to recover a pure condensate.
- [19] C. C. Bradley, C. A. Sackett, and R. G. Hulet, Phys. Rev. Lett. **78**, 985 (1997).
- [20] The *in situ* images are taken immediately (within 5 μs) after the cloud is released from the optical trap.
- [21] D. Clément, P. Bouyer, A. Aspect, and L. Sanchez-Palencia, preceding paper, Phys. Rev. A **77**, 033631 (2008).
- [22] S. Dettmer *et al.*, Phys. Rev. Lett. **87**, 160406 (2001).
- [23] Z. Hadzibabic, S. Stock, B. Battelier, V. Bretin, and J. Dalibard, Phys. Rev. Lett. **93**, 180403 (2004).
- [24] L. Fallani, C. Fort, J. E. Lye, and M. Inguscio, Opt. Express **13**, 4303 (2005).
- [25] L. Sanchez-Palencia, D. Clément, P. Lugan, P. Bouyer, G. V. Shlyapnikov, and A. Aspect, Phys. Rev. Lett. **98**, 210401 (2007); L. Sanchez-Palencia, Phys. Rev. A **74**, 053625 (2006); P. Lugan, D. Clément, P. Bouyer, A. Aspect, M. Lewenstein, and L. Sanchez-Palencia, Phys. Rev. Lett. **98**, 170403 (2007).
- [26] G. Roati, M. Zaccanti, C. D'Errico, J. Catani, M. Modugno, A. Simoni, M. Inguscio, and G. Modugno, Phys. Rev. Lett. **99**, 010403 (2007).
- [27] T. Paul, P. Schlagheck, P. Leboeuf, and N. Pavloff, Phys. Rev. Lett. **98**, 210602 (2007).
- [28] P. W. Anderson, *Basic Notions of Condensed Matter Physics* (Westview, Boulder, CO, 1997).
- [29] D. McKay, M. White, M. Pasienski, and B. DeMarco, e-print arXiv:0708.3074.