

The Postulates Of Quantum Mechanics

Postulate 1: The State of the System

The set of all possible states of a physical system stand in a one-to-one correspondence with the vector directions (rays) in a Hilbert space \mathcal{H} . (\mathcal{H} can be finite or infinite depending upon the system. To allow for continuous basis vectors we will extend the system and space to $\hat{\mathcal{H}}$.)

Since the state is described by the entire ray, we have that $|\psi\rangle$ and $\lambda|\psi\rangle$ with $\lambda \in \mathcal{C}$ describe the same state.

Postulate 2: Physical Observables and Hermitian Operators

The physical observables of a system stand in a one-to-one correspondence with the set of Hermitian operators on the state space \mathcal{H} . That is, to each measurable quantity \mathcal{A} , there corresponds a Hermitian operator A acting on \mathcal{H} . Out of the set of all Hermitian operators, there is a subset which consists of mutually commuting operators and are assumed to be complete (they form a CSCO; each A is assumed an observable).

Postulate 3: Measurement, Spectral Decomposition and Statistical Interpretation

- a. The only possible result of the measurement of a physical observable \mathcal{A} is one of the (real) eigenvalues of the corresponding Hermitian operator A .
- b. Let $\{|\phi_k\rangle\}$ be the simultaneous eigenstates of a CSCO so that

$$A|\phi_k\rangle = a_k|\phi_k\rangle, \quad \text{etc.} \quad (1)$$

These states form an orthonormal basis for the state space \mathcal{H} .

- c. For a system in state $|\psi\rangle$ (with $\langle\psi|\psi\rangle = 1$) the probability of measuring the value a_k for the physical observable \mathcal{A} is

$$P_k = |\langle\phi_k|\psi\rangle|^2. \quad (2)$$

d. Immediately following the measurement, the system is in the state $|\phi_k\rangle$.

To be more explicit:

Discrete Orthonormal Basis

If the orthonormal basis is a discrete basis, then the simultaneous eigenvectors, $|\phi_{ab\dots}\rangle$, of the CSCO $\{A, B, \dots\}$ obey

$$\begin{aligned} \langle \phi_{a'b'\dots} | \phi_{ab\dots} \rangle &= \delta_{a'a} \delta_{b'b} \dots \\ \sum_{a,b,\dots} |\phi_{ab\dots}\rangle \langle \phi_{ab\dots}| &= 1, \end{aligned} \quad (3)$$

where

$$\begin{aligned} A|\phi_{ab\dots}\rangle &= a|\phi_{ab\dots}\rangle \\ B|\phi_{ab\dots}\rangle &= b|\phi_{ab\dots}\rangle, \quad \text{etc.}, \end{aligned} \quad (4)$$

with the eigenvalues $\{a, b, \dots\}$ taking discrete values (in 1-1 correspondence with the integers).

For an arbitrary state vector $|\psi\rangle$, we have the expansion in terms of the $\{|\phi_{ab\dots}\rangle\}$ basis

$$|\psi\rangle = \sum_{a,b,\dots} \psi_{ab\dots} |\phi_{ab\dots}\rangle, \quad (5)$$

with

$$\psi_{ab\dots} = \langle \phi_{ab\dots} | \psi \rangle. \quad (6)$$

This implies

$$\begin{aligned} A|\psi\rangle &= \sum_{a,b,\dots} a \psi_{ab\dots} |\phi_{ab\dots}\rangle \\ B|\psi\rangle &= \sum_{a,b,\dots} b \psi_{ab\dots} |\phi_{ab\dots}\rangle, \\ \text{etc.} & \end{aligned} \quad (7)$$

The probability of finding the values a, b, \dots for the system in state $|\psi\rangle$ when A, B, \dots are measured is

$$P_{ab\dots} = |\langle \phi_{ab\dots} | \psi \rangle|^2. \quad (8)$$

Note that

$$\begin{aligned}
 \sum_{a,b,\dots} P_{ab\dots} &= \sum_{a,b,\dots} |\langle \phi_{ab\dots} | \psi \rangle|^2 \\
 &= \sum_{a,b,\dots} \langle \psi | \phi_{ab\dots} \rangle \langle \phi_{ab\dots} | \psi \rangle \\
 &= \langle \psi | \underbrace{\sum_{a,b,\dots} |\phi_{ab\dots} \rangle \langle \phi_{ab\dots} |}_{=1} | \psi \rangle \\
 &= \langle \psi | \psi \rangle \\
 &= 1, \tag{9}
 \end{aligned}$$

as required of a probability. Also the expectation value of A, B, \dots in state $|\psi\rangle$ is

$$\begin{aligned}
 \langle \psi | A | \psi \rangle &= \sum_{a,b,\dots} a \underbrace{\langle \psi | \phi_{ab\dots} \rangle}_{=\langle \phi_{ab\dots} | \psi \rangle} \\
 &= \sum_{a,b,\dots} a |\langle \phi_{ab\dots} | \psi \rangle|^2 \\
 &= \sum_{a,b,\dots} a P_{ab\dots}, \quad \text{etc.} \tag{10}
 \end{aligned}$$

Continuous Orthonormal Basis

On the otherhand if the orthonormal basis is continuous, then

$$\begin{aligned}
 \langle \phi_{\alpha'\beta'\dots} | \phi_{\alpha\beta\dots} \rangle &= \delta(\alpha' - \alpha) \delta(\beta' - \beta) \dots \\
 \int d\alpha d\beta \dots |\phi_{\alpha\beta\dots} \rangle \langle \phi_{\alpha\beta\dots} | &= 1, \tag{11}
 \end{aligned}$$

where

$$\begin{aligned}
 A |\phi_{\alpha\beta\dots} \rangle &= \alpha |\phi_{\alpha\beta\dots} \rangle \\
 B |\phi_{\alpha\beta\dots} \rangle &= \beta |\phi_{\alpha\beta\dots} \rangle, \\
 &\text{etc.} \tag{12}
 \end{aligned}$$

with the eigenvalues $\{\alpha, \beta, \dots\}$ taking on a continuum of values.

For an arbitrary ket vector $|\psi\rangle$, the expansion in terms of the continuous basis $\{|\phi_{\alpha\beta\dots}\rangle\}$ is

$$|\psi\rangle = \int d\alpha d\beta \dots \psi(\alpha, \beta, \dots) |\phi_{\alpha\beta\dots}\rangle, \tag{13}$$

with

$$\psi(\alpha, \beta, \dots) = \langle \phi_{\alpha\beta\dots} | \psi \rangle . \quad (14)$$

This implies

$$\begin{aligned} A|\psi\rangle &= \int d\alpha d\beta \dots \alpha \psi(\alpha, \beta, \dots) |\phi_{\alpha\beta\dots}\rangle \\ B|\psi\rangle &= \int d\alpha d\beta \dots \beta \psi(\alpha, \beta, \dots) |\phi_{\alpha\beta\dots}\rangle \\ \text{etc.} & \end{aligned} \quad (15)$$

The probability of measuring \mathcal{A} in the range α to $\alpha + d\alpha$, \mathcal{B} in the range β to $\beta + d\beta$, etc. is

$$d\mathcal{P}(\alpha, \beta, \dots) = |\langle \phi_{\alpha\beta\dots} | \psi \rangle|^2 d\alpha d\beta \dots . \quad (16)$$

Note that

$$\begin{aligned} \int d\mathcal{P}(\alpha, \beta, \dots) &= \int d\alpha d\beta \dots |\langle \phi_{\alpha\beta\dots} | \psi \rangle|^2 \\ &= \langle \psi | \underbrace{\int d\alpha d\beta \dots |\phi_{\alpha\beta\dots}\rangle \langle \phi_{\alpha\beta\dots}|}_{=1} | \psi \rangle \\ &= \langle \psi | \psi \rangle \\ &= 1 , \end{aligned} \quad (17)$$

as required of a probability. Also the expectation value of A, B, \dots in state $|\psi\rangle$ is

$$\begin{aligned} \langle \psi | A | \psi \rangle &= \int d\alpha d\beta \dots \alpha \underbrace{\psi(\alpha, \beta, \dots)}_{=\langle \phi_{\alpha\beta\dots} | \psi \rangle} \langle \psi | \phi_{\alpha\beta\dots} \rangle \\ &= \int d\alpha d\beta \dots \alpha |\langle \phi_{\alpha\beta\dots} | \psi \rangle|^2 \\ &= \int d\mathcal{P}(\alpha, \beta, \dots) \alpha \\ \text{etc.} & \end{aligned} \quad (18)$$

Postulate 4: Time Evolution and The Schrödinger Equation

The time evolution of the physical states is given by the Schrödinger equation

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H(t) |\psi(t)\rangle , \quad (19)$$

where $H = H(t)$ is the Hermitian Hamiltonian operator. $H(t)$ is the total energy of the system.

For an isolated system, the Hamiltonian is time independent. Schrödinger's equation is simply

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H |\psi(t)\rangle, \quad (20)$$

with the eigenvalues of H the possible energies of the system.

In general the observables may also have explicit time dependence $A = A(t)$. (For example, a charged particle can interact with an external time varying electromagnetic field.) For an isolated system, as is the Hamiltonian, the observables are time independent, $\frac{dA}{dt} = 0$.