

Physics 22000 General Physics

Lecture 25 – Waves

Fall 2016 Semester

Prof. Matthew Jones

1

Final Exam

UNITIME Examination Schedule Fall 2016 (PWL) [Click here to change the semester/term](#)

Filter
Term: Fall 2016 (PWL) Exams: Final Subject: PHYS Apply

Fall 2016 (PWL) final examinations (PHYS)

Subject	Course	CRN	Section	Date	Time	Room
PHYS	17200			Wed 12/14	1:00p - 3:00p	LAMB F101
PHYS	17200H	26979	26979-H01	Mon 12/12	8:00a - 10:00a	PHYS 114
PHYS	21400	26984	26984-001	Tue 12/13	10:30a - 12:30p	PHYS 114
PHYS	21500	26987	26987-001	Fri 12/16	1:00p - 3:00p	PHYS 112
PHYS	21800	16840	16840-014	Mon 12/12	7:00p - 9:00p	HKRS B848
PHYS	21800	26997	26997-001	Fri 12/16	1:00p - 3:00p	PHYS 114
PHYS	21900	27009	27009-001	Wed 12/14	1:00p - 3:00p	PHYS 112
PHYS	22000			Fri 12/16	1:00p - 3:00p	LAMB F101
PHYS	22100			Wed 12/14	1:00p - 3:00p	LILY 1105
PHYS	23300	67161	67161-001	Fri 12/16	8:00a - 10:00a	EE 129
PHYS	23400	10833	10833-001	Fri 12/16	8:00a - 10:00a	PHYS 111

2

Free Study Sessions!

Rachel Hoagburg

Come to SI for more help in **PHYS 220**

Tuesday and Thursday 7:30-8:30PM Shreve C113

Office Hour


Tuesday 1:30-2:30 4th floor of Krach

For other SI-linked courses and schedules, visit purdue.edu/si or purdue.edu/bolleguide

3

Mechanical Waves

- Waves and wave fronts:




Wave Motion

Wave motion involves a disturbance produced by a vibrating object (a source). The disturbance moves, or propagates, through a medium and causes points in the medium to vibrate. When the disturbed medium is physical matter (solid, liquid, or gas), the wave is called a **mechanical wave**.


Two Kinds of Waves

Longitudinal and transverse waves In a *longitudinal wave* the vibrational motion of the particles or layers of the medium is parallel to the direction of propagation of the disturbance. In a *transverse wave* the vibrational motion of the particles or layers of the medium is perpendicular to the direction of propagation of the disturbance.



Reflection of Waves

- When a wave reaches the wall of the container or the end of the Slinky or rope, it reflects off the end and moves in the opposite direction.
 - When a wave encounters any boundary between different media, some of the wave is reflected back.




7

Mathematical Description of Waves

- A wave can be created in a rope by a motor that vibrates the end of a rope up and down, producing a transverse wave.
- The displacement is described by a sinusoidal function of time:

$$y = A \cos\left(\frac{2\pi}{T} t\right)$$



8

Mathematical Description of Waves

Period T in seconds is the time interval for one complete vibration of a point in the medium anywhere along the wave's path.

Frequency f in Hz (s^{-1}) is the number of vibrations per second of a point in the medium as the wave passes.

Amplitude A is the maximum displacement of a point of the medium from its equilibrium position as the wave passes.

Speed v in m/s is the distance a disturbance travels during a time interval, divided by that time interval.

9

Mathematical Description of a Traveling Sinusoidal Wave

- We know the source oscillates up and down with a vertical displacement given by:

$$y = A \cos\left(\frac{2\pi}{T} t\right)$$

- We can mathematically describe the disturbance $y(x, t)$ of a point of the rope at some positive position x to the right of the source (at $x = 0$) by:

$$y(x, t) = A \cos\left[\frac{2\pi}{T} \left(t - \frac{x}{v}\right)\right]$$

10

Wavelength

Wavelength λ equals the distance between two nearest points on a wave that at any clock reading have exactly the same displacement and shape (slope). It is also the distance between two consecutive wave fronts:

$$\lambda = Tv = \frac{v}{f} \quad (20.3)$$

11

Mathematical Description of a Traveling Sinusoidal Wave

Mathematical description of a traveling sinusoidal wave The displacement from equilibrium y of a point at location x at time t when a wave of period T travels at speed v in the positive x -direction through a medium is described by the function

$$y = A \cos\left[2\pi\left(\frac{t}{T} - \frac{x}{\lambda}\right)\right] \quad (20.4)$$

The wavelength λ of this wave equals $\lambda = Tv$.

12

Wave Speed

- The speed of a wave on a string depends on
 - The mass per unit length (μ): waves propagate more quickly on a light string
 - The tension (T): waves propagate more quickly when the string is tight


$$v = \sqrt{\frac{T}{\mu}}$$

13


Amplitude and Energy in a Two Dimensional Medium

- A beach ball bobs up and down in water in simple harmonic motion, producing circular waves that travel outward across the water surface in all directions.
 - The amplitudes of the crests decrease as the waves move farther from the source.

Top view of wave crests at one instant in time



Side view of wave crests at one instant in time




14

Amplitude and Energy in a Two Dimensional Medium

- The circumference of the second ring is two times greater than the first, but the same energy per unit time moves through it.
 - The energy per unit circumference length passing through the second ring is one-half that passing through the first ring.

Snapshot of wave crests at one instant in time



The same energy/time passes two rings that have different circumferences.

15


Two-dimensional waves produced by a point source

Two-dimensional waves produced by a point source The energy per unit circumference length and per unit time crossing a line perpendicular to the direction that the wave travels decreases as $1/r$, where r is the distance from the point source of the wave.

16

Three-Dimensional waves produced by a point source

- The area of the second sphere is four times the area of the first sphere, but the same energy per unit time moves through it.
 - The energy per unit area through the second sphere is one-fourth that through the first sphere.



17

Three-dimensional waves produced by a point source

Three-dimensional waves produced by a point source The energy per unit area per unit time passing across a surface perpendicular to the direction that the wave travels decreases as $1/r^2$, where r is the distance from the point source of the wave.

18

Wave Power and Wave Intensity

- The intensity of a wave is defined as the energy per unit area per unit time interval that crosses perpendicular to an area in the medium through which it travels:

$$\text{Intensity} = \frac{\text{Energy}}{\text{Time} \cdot \text{Area}} = I = \frac{\Delta U}{\Delta t \cdot A} = \frac{P}{A}$$

- The unit of intensity I is equivalent to joules per second per square meter or watts per square meter.


19

Reflection and Impedance

- If you hold one end of a rope whose other end is fixed and shake it once, you create a transverse traveling incident pulse.
 - When the pulse reaches the fixed end, the reflected pulse bounces back in the opposite direction.
 - The reflected pulse is inverted—oriented downward as opposed to upward.
- What happens to a wave when there is an abrupt change from one medium to another?

20

Reflection and Impedance



21

Impedance

- Impedance characterizes the degree to which waves are reflected and transmitted at the boundary between different media.
- Impedance is defined as the square root of the product of the elastic and inertial properties of the medium:

$$\text{Impedance} = Z = \sqrt{(\text{Elastic property})(\text{Inertial property})}$$

22

Amplitude of Reflected Waves

- The amplitude of the reflected pulse depends on the impedance of the two media:

$$A_r = R A_i$$

$$R = \frac{Z_2 - Z_1}{Z_2 + Z_1}$$

(when incident from medium 1 into medium 2)



23

Ultrasound

- Ultrasound takes advantage of the differing densities of internal structures to "see" inside the body.
 - The impedance of tissue is much greater than that of air. As a result, most of the ultrasound energy is reflected at the air-body interface and does not travel inside the body.
 - To overcome this problem, the area of the body to be scanned is covered with a gel that helps "match" the impedance between the emitter and the body surface.

24


Superposition Principle

- We have studied the behavior of a single wave traveling in a medium and initiated by a simple harmonic oscillator.
 - Most periodic or repetitive disturbances of a medium are combinations of two or more waves of the same or different frequencies traveling through the same medium at the same time.
- Now we investigate what happens when two or more waves simultaneously pass through a medium.

25

Superposition Principle


- Imagine we have two vibrating sources in water, each of which sends out sinusoidal waves.
 - Consider point C in the figure.



26

Superposition Principle

- Imagine we have two vibrating sources in water, each of which sends out sinusoidal waves.
 - Consider point D in the figure.




Superposition principle for waves

- The process in which two or more waves of the same frequency overlap is called interference.
 - Places where the waves add to create a larger disturbance are called locations of constructive interference.
 - Places where the waves add to produce a smaller disturbance are called locations of destructive interference.

28


Huygens' Principle

- We can explain the formation of the waves using the superposition principle, first explained by Christiaan Huygens (1629–1695).
 - Each arc represents a wavelet that moves away from a point on the original wave front.



29

Huygens' Principle



30

Loudness and intensity

- Loudness is determined primarily by the amplitude of the sound wave: the larger the amplitude, the louder the sound.
 - Equal-amplitude sound waves of different frequencies will not have the same perceived loudness to humans.
- To measure the loudness of a sound, we measure the intensity: the energy per unit area per unit time interval.

31

Intensity Level

- Because of the wide variation in the range of sound intensities, a quantity called intensity level is commonly used to compare the intensity of one sound to the intensity of a reference sound.
- Intensity level is defined on a base 10 logarithmic scale as follows:

$$\beta = \log_{10} \frac{I}{I_0}$$

32

Intensities and Intensity Levels of Common Sounds

Table 20.6 Intensities and intensity levels of common sounds.

Source of sound	Intensity (W/m ²)	Intensity level (dB)	Description
Large rocket engine (nearby)	10 ⁶	180	
Jet takeoff (nearby)	10 ³	150	
Pneumatic riveter; machine gun (nearby)	10	130	
Rock concert with amplifiers (2 m); jet takeoff (60 m)	1	120	Pain threshold
Construction noise (3 m)	10 ⁻¹	110	
Moving subway train (nearby)	10 ⁻²	100	
Heavy truck (15 m)	10 ⁻³	90	Constant exposure endangers hearing
Niagara Falls (nearby)	10 ⁻³	90	
Noisy office with machines; inside an average factory	10 ⁻⁴	80	
Busy traffic	10 ⁻⁵	70	
Normal conversation (1 m)	10 ⁻⁶	60	
Quiet office	10 ⁻⁷	50	Quiet
Library	10 ⁻⁸	40	
Soft whisper (5 m)	10 ⁻⁹	30	
Rustling leaves	10 ⁻¹⁰	20	Barely audible
Normal breathing	10 ⁻¹¹	10	
	10 ⁻¹²	0	Hearing threshold

33


Pitch, frequency and complex sounds

- Pitch is the perception of the frequency of a sound.
 - Tuning forks of different sizes produce sounds of approximately the same intensity, but we hear each as having a different pitch.
 - The shorter the length of the tuning fork, the higher the frequency and the higher the pitch.
- Like loudness, pitch is not a physical quantity but rather a subjective impression.

34

Complex sounds, waveforms and frequency spectra


- An oboe and a violin playing concert A equally loudly at 440 Hz sound very different.
 - Which other property of sound causes this different sensation to our ears?
 - Musical notes from instruments have a characteristic frequency, but the wave is not sinusoidal.



35

Beat and Beat Frequencies

- Two sound sources of similar (but not the same) frequency are equidistant from a microphone that records the air pressure variations due to the two sound sources as a function of time.



36

Beat and beat frequencies


A **beat** is a wave that results from the superposition of two waves of about the same frequency. The beat (the net wave) has a frequency equal to the average of the two frequencies and has variable amplitude. The frequency with which the amplitude of the net wave changes is called the **beat frequency** f_{beat} ; it equals the difference in the frequencies of the two waves:

$$f_{\text{beat}} = |f_1 - f_2| \quad (20.10)$$

37

Standing Waves on Strings

- You shake the end of a rope that is attached to a fixed support.
 - At specific frequencies, you notice that the rope has large-amplitude sine-shaped vibrations that appear not to be traveling.




38

Standing Waves on a String

- The lowest-frequency vibration of the rope is one up-and-down shake per time interval:

$$f_1 = \frac{1}{\tau} = \frac{1}{2L/v} = \frac{v}{2L}$$

- This frequency is called the **fundamental frequency**.



39
