




Fireworks from Magnetar Birth

Gamma-Ray Bursts
(Long and Short)



Super-Luminous
Supernovae



Brian Metzger
Columbia University

In Collaboration with

Indrek Vurm, Romain Hascoet, Andrei Beloborodov (Columbia), Tony Piro (Caltech)
Eliot Quataert, Geoff Bower, Jon Arons (UC Berkeley)
Niccolo Bucciantini (INAF), Todd Thompson (OSU), Dimitrios Giannios (Purdue)
Paul O' Brien, A. Levan (Leicester), A. Rowlinson (Amsterdam)

GRB Magnetar Thinkshop

Bormio, Italy - January 21, 2014

‘Millisecond Magnetars’

All neutron stars form as hot,
differentially-rotating ‘proto-neutron stars’

rotational
energy:

The image cannot be displayed. Your computer may not have enough memory to open the image, or the image may have been corrupted. Restart your computer, and then open the file again. If the red x still appears, you may have to delete the image and then insert it again.


The image cannot be displayed. Your computer may not have enough memory to open the image, or the image may have been corrupted. Restart your computer, and then open the file again. If the red x still appears, you may have to delete the image and then insert it again.

Field amplification:

- Shear instabilities (talk by Zrake)
- Magneto-rotational instability
- α - Ω dynamo (Thompson & Duncan 1993)

The image cannot be displayed. Your computer may not have enough memory to open the image, or the image may have been corrupted. Restart your computer, and then open the file again. If the red x still appears, you may have to delete the image and then insert it again.

$Ro \sim 1$ for $P \sim 1$ ms



Magnetic activity of late type stars

Rossby Number $\text{Log}(Ro = P/\tau_c)$

Pizzolato et al. 2003



Signatures of Magnetar Birth

spin-down .
luminosity :

$$L_{\text{sd}} = \frac{\mu^2 \Omega^4}{c^3} \approx 6 \times 10^{49} \left(\frac{P}{1 \text{ ms}} \right)^{-4} \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^2 \text{ erg s}^{-1}$$

spin-down time :

$$\tau_{\text{sd}} = \frac{E_{\text{rot}}}{L_{\text{sd}}} \approx 10 \left(\frac{P_0}{1 \text{ ms}} \right)^2 \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^{-2} \text{ min}$$




Signatures of Magnetar Birth

spin-down luminosity :

$$L_{\text{sd}} = \frac{\mu^2 \Omega^4}{c^3} \approx 6 \times 10^{49} \left(\frac{P}{1 \text{ ms}} \right)^{-4} \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^2 \text{ erg s}^{-1}$$


spin-down time :

$$\tau_{\text{sd}} = \frac{E_{\text{rot}}}{L_{\text{sd}}} \approx 10 \left(\frac{P_0}{1 \text{ ms}} \right)^2 \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^{-2} \text{ min}$$



Gamma-Ray Burst

- Jet punches successfully through star
- $L_{\text{sd}} \sim L_{\gamma} \sim 10^{49-51} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{minutes-hours}$




Signatures of Magnetar Birth

spin-down luminosity :

$$L_{\text{sd}} = \frac{\mu^2 \Omega^4}{c^3} \approx 6 \times 10^{49} \left(\frac{P}{1 \text{ ms}} \right)^{-4} \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^2 \text{ erg s}^{-1}$$

spin-down time :

$$\tau_{\text{sd}} = \frac{E_{\text{rot}}}{L_{\text{sd}}} \approx 10 \left(\frac{P_0}{1 \text{ ms}} \right)^2 \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^{-2} \text{ min}$$




Gamma-Ray Burst

- Jet punches successfully through star
- $L_{\text{sd}} \sim L_{\gamma} \sim 10^{49-51} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{minutes-hours}$

Super-Luminous SN

- Jet stifled, but optical SN powered diffusively
- $L_{\text{sd}} \sim L_{\text{SN}} \sim 10^{43-45} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{week - months}$




Signatures of Magnetar Birth

spin-down luminosity :

$$L_{\text{sd}} = \frac{\mu^2 \Omega^4}{c^3} \approx 6 \times 10^{49} \left(\frac{P}{1 \text{ ms}} \right)^{-4} \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^2 \text{ erg s}^{-1}$$

spin-down time :

$$\tau_{\text{sd}} = \frac{E_{\text{rot}}}{L_{\text{sd}}} \approx 10 \left(\frac{P_0}{1 \text{ ms}} \right)^2 \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^{-2} \text{ min}$$




Gamma-Ray Burst

- Jet punches successfully through star
- $L_{\text{sd}} \sim L_{\gamma} \sim 10^{49-51} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{minutes-hours}$


Super-Luminous SN

- Jet stifled, but optical SN powered diffusively
- $L_{\text{sd}} \sim L_{\text{SN}} \sim 10^{43-45} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{week - months}$



Constraints on the GRB Central Engine

- Energies - $E_\gamma \sim 10^{49-52}$ ergs
- Rapid ~ms variability
- Duration - $T_\gamma \sim 10-100$ s
- Steep decay phase
- Narrowly collimated jet
- Bulk Lorentz factor $\Gamma \sim 100-1000$ ($M_{jet} < 10^{-5} M_\odot$)
- Late activity (plateau & flaring)



BH

versus

NS

GAMMA-RAY BURSTS FROM STELLAR MASS ACCRETION DISKS AROUND BLACK HOLES¹


S. E. WOOSLEY

University of California Observatories/Lick Observatory, Board of Studies in Astronomy and Astrophysics, University of California, Santa Cruz, Santa Cruz, CA 95064; and General Studies Group, Physics Department, Lawrence Livermore National Laboratory

Received 1992 June 22; accepted 1992 September 3


ABSTRACT

A cosmological model for gamma-ray bursts is explored in which the radiation is produced as a broadly beamed pair fireball along the rotation axis of an accreting black hole. The black hole may be a consequence of neutron star merger or neutron star–black hole merger, but for long complex bursts, it is more likely to come from the collapse of a single Wolf-Rayet star endowed with rotation (“failed” Type Ib supernova). The





GRB SNe are
actually quite
successful!


Bright \Rightarrow
 $M_{Ni56} > 0.1 M_\odot$
Energetic $\Rightarrow E_{KE}$
 $\sim 10^{52}$ ergs



...but massive stars (ZAMS $>25 M_{\odot}$)
become black holes, right?









Core Collapse with Magnetic Fields & Rotation

(e.g. LeBlanc & Wilson 1970)


THE PROTO-NEUTRON STAR PHASE OF THE COLLAPSAR MODEL AND THE ROUTE TO LONG-SOFT GAMMA-RAY BURSTS AND HYPERNOVAE

L. DESSART¹, A. BURROWS¹, E. LIVNE², AND C.D. OTT¹




THE ARDUOUS JOURNEY TO BLACK-HOLE FORMATION IN POTENTIAL GAMMA-RAY BURST PROGENITORS

LUC DESSART,^{1,2} EVAN O'CONNOR,² AND CHRISTIAN D. OTT^{2,3,*}



Alternative View of the Fates of Massive *Rotating* Stars

(Metzger et al. 2011; see also Dessart, O' Connor, & Ott 2012)



Neutrino Driven Wind


Neutrinos heat proto-NS atmosphere (e.g. $\nu_e + n \Rightarrow p + e^-$)
⇒ **drives wind behind outgoing supernova shock** (e.g. Qian & Woosley 96)



$$\dot{M} \sim 10^{-4} \left(\frac{L_\nu}{10^{52} \text{erg s}^{-1}} \right)^{5/3} \left(\frac{\varepsilon_\nu}{10 \text{ MeV}} \right)^{10/3} M_\odot \text{s}^{-1} \Rightarrow \text{crucial to baryon loading}$$

Effects of Strong Magnetic Fields


“Helmet - Streamer”



- Microphysics (EOS, ∇ Heating & Cooling)
 - Important for $B \geq 10^{16}$ G (Duan & Qian 2005)

Effects of Strong Magnetic Fields

“Helmet - Streamer”




- Microphysics (EOS, ν Heating & Cooling)
 - Important for $B \geq 10^{16}$ G (Duan & Qian 2005)
- Magneto-Centrifugal Slinging
(Weber & Davis 1967; Thompson, Chang & Quataert 2004)

Outflow Co-Rotates
with Neutron Star when

$$\frac{B^2}{8\pi} > \frac{1}{2} \rho v_r^2$$


⇒

Magneto-Centrifugal
Acceleration (“Beads
on a Wire”)




Regimes of Magnetized PNS Winds ($B = 3 \times 10^{14}$ G)

Metzger, Thompson, Quataert 2007





Regimes of Magnetized PNS Winds ($B = 3 \times 10^{14}$ G)

Metzger, Thompson, Quataert 2007



Evolution of Proto-Magnetar Outflows

(BDM et al. 2007, 2011)




Wind Power $\dot{E}(t)$, Mass Loss Rate $\dot{M}(t)$,

Calculate:

$$\Rightarrow \text{'Magnetization'} \ \sigma(t) \sim \dot{E} / \dot{M} c^2 = \Gamma_{\max}(t)$$

In terms of

Initial rotation period P_0 , **dipole field B_{dip}** & **obliquity θ_{dip}**





$$\sigma_0 \sim \Gamma_{\max} = \frac{\dot{E}}{\dot{M}c^2} \propto \frac{B^2 \Omega^4}{L_\nu^{5/3} T^{10/3}}$$

increases as magnetar cools



Collimation via Stellar Confinement

Multi-Wavelength Crab Nebula




Collimation via Stellar Confinement

Multi-Wavelength Crab Nebula




Supernova remnant elongated by
anisotropic magnetic stresses
in pulsar nebula? (Begelman & Li 1992)




Stellar Mantle


Outgoing SN shock
 $V_{SN} \sim 0.1 c$




Outgoing SN shock
 $V_{SN} \sim 0.1 c$



Outgoing SN shock
 $V_{SN} \sim 0.1 c$





Outgoing SN shock
 $V_{SN} \sim 0.1 c$




Jet Formation via Stellar Confinement


(Bucciantini et al. 2007, 08, 09; cf. Uzdensky & MacFadyen 07; Komissarov & Barkov 08)




Kink Instability



Porth, Komissarov, & Keppens 13




Outflow becomes relativistic at $t \sim 2$ seconds;
Jet breaks out of star at $t_{\text{bo}} \sim R_\star/\beta c \sim 10$ seconds




Outflow becomes relativistic at $t \sim 2$ seconds;
Jet breaks out of star at $t_{\text{bo}} \sim R_\star/\beta c \sim 10$ seconds


GRB Emission - What, Where, How?



1. **What** is jet's composition? (kinetic or magnetic?)
2. **Where** is dissipation occurring? (photosphere? deceleration radius?)
3. **How** is radiation generated? (synchrotron, IC, hadronic?)


Metzger et al. 2011






Metzger et al. 2011


Hot Electrons \Rightarrow
IC Scattering (γ -rays)
and Synchrotron (optical)



E_{peak} Evolution



Ave. Magnetization σ_{avg}



$\sigma_{\text{avg}}\text{-}L_\gamma$ Correlation

Prediction:
More Luminous GRBs
 \Leftrightarrow Higher Γ

Ave. Wind Power (erg s^{-1})

Consistent with

$E_{\text{peak}} \propto E_{\text{iso}}^{0.4}$ (Amati +02)


$E_{\text{peak}} \propto L_{\text{iso}}^{0.5}$

(Yonetoku+04)


And $\Gamma \propto E_{\text{iso}}^{0.3}$

(Liang+10)

Average E_{peak} (keV)



End of the GRB = Neutrino Transparency?



Ultra High- σ Outflows

\Rightarrow

- Acceleration is Inefficient

(e.g. Tchekhovskoy et al. 2009)

- Internal Shocks are Weak


(e.g. Kennel & Coroniti 1984)

- Reconnection is Slow

(e.g. Drenkahn & Spruit 2002)


$$T_{\text{GRB}} \sim T_{\nu \text{ thin}} \sim 20 - 100 \text{ s}$$

End of the GRB = Neutrino Transparency?




Ultra High- σ Outflows
 ⇒


- Acceleration is Inefficient
 (e.g. Tchekhovskoy et al. 2009)
- Internal Shocks are Weak
 (e.g. Kennel & Coroniti 1984)
- Reconnection is Slow
 (e.g. Drenkahn & Spruit 2002)



$$T_{\text{GRB}} \sim T_{\nu \text{ thin}} \sim 20 - 100 \text{ s}$$




e.g. Zhang & Meszaros 2001; Troja et al. 2007; Yu et al. 2009; Lyons et al. 2010




Plateau Duration - Luminosity Correlation

Spin-Down Timescale




‘Plateau’ Luminosity


A Diversity of Magnetar Birth



A Diversity of Magnetar Birth




A Diversity of Magnetar Birth



Observational Tests & Constraints

- Max Energy* -
 $E_{KE} + E_\gamma < 3 \times 10^{52}$ ergs
- *subject to uncertainties in afterglow modeling.
(e.g. Zhang & MacFadyen 09).



- Long GRB *always* accompanied by bright, energetic
 - Consistent with observations thus far (Woosley & Bloom 2006).
- Γ increases during GRB and correlates with E_γ
 - translate jet luminosity/magnetization into unique prediction for gamma-ray light curves and spectra.


Signatures of Magnetar Birth

spin-down luminosity :

$$L_{\text{sd}} = \frac{\mu^2 \Omega^4}{c^3} \approx 6 \times 10^{49} \left(\frac{P}{1 \text{ ms}} \right)^{-4} \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^2 \text{ erg s}^{-1}$$

spin-down time :

$$\tau_{\text{sd}} = \frac{E_{\text{rot}}}{L_{\text{sd}}} \approx 10 \left(\frac{P_0}{1 \text{ ms}} \right)^2 \left(\frac{B_{\text{dip}}}{10^{15} \text{ G}} \right)^{-2} \text{ min}$$




Gamma-Ray Burst

- Jet punches successfully through star
- $L_{\text{sd}} \sim L_{\gamma} \sim 10^{49-51} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{minutes-hours}$

Super-Luminous SN

- Jet stifled, but optical SN powered diffusively
- $L_{\text{sd}} \sim L_{\text{SN}} \sim 10^{43-45} \text{ erg s}^{-1}$
- $\tau_{\text{sd}} \sim \text{week - months}$



Summary of the Proto-Magnetar Model for GRBs

- ✓ **GRB Duration $\sim 10 - 100$ seconds & Steep Decay Phase**

- Time for NS to become transparent to neutrinos (end of ν-wind)

- ✓ **GRB Energies $E_{\text{GRB}} \sim 10^{50-52}$ ergs**

- Rotational energy lost in $\sim 10-100$ s

- ✓ **Ultra-Relativistic Outflow with $\Gamma \sim 100-1000$**

- Mass loading set by physics of neutrino heating (not fine-tuned).

- ✓ **Jet Collimation**

- Star confines and redirects magnetar outflow into jet

- ✓ **Association with Energetic Core Collapse Supernovae**

- $E_{\text{rot}} \sim E_{\text{SN}} \sim 10^{52}$ ergs - MHD-powered SN associated w magnetar birth.



- ✓ **Late-Time Central Engine Activity**

- Residual rotational (plateau) or magnetic energy (flares)?

Hydrogen-Poor ‘SuperLuminous’ Supernovae

Quimby+07, Barbary+09, Pastorello+10, Chomiuk+11, Leloudas+12, Berger+12, Lunnan+13, Inserra+13; Nicholl+13; McCrum+14

Quimby et al. 2011





- $L_{\text{peak}} > 10^{44} \text{ erg s}^{-1}$, $E_{\text{rad}} \sim 10^{50-51} \text{ ergs}$ ($10-100 \times$ normal SNe)
- UV-rich spectrum with intermediate mass elements
- Faint metal-poor host galaxies, similar to long GRBs
(Quimby+11, Neill+11, Chomiuk+11, Chen+13; but see Berger+13, Chornock+13)
- Competing models: circumstellar interaction vs. central engine

Millisecond Magnetar-Powered Supernovae

(Kasen & Bildsten 2010; Woosley 2010; Dessart et al. 2011)

Kasen & Bildsten 2010



PROS


- SN luminosity increased if pulsar spin-down time \sim optical peak t_{peak} $\Rightarrow B_{\text{dip}} \sim 10^{13-14}$ G
- Explains similar host galaxies to long GRBs (both require rapidly rotating progenitor)
- Can reproduce diversity of rise times and peak luminosities

CONS


- Can reproduce diversity of rise times and peak luminosities (hard to test)
 - Difficult to distinguish from other ‘hidden’ energy sources (optically-thick CSM interaction)
 - Assumes pulsar luminosity thermalized
- Reality: Poynting flux $\Rightarrow e^{+/-} \Rightarrow$ non-thermal radiation \Rightarrow thermal radiation**

How Supernovae Shine (Arnett 1982)

SN ejecta w mass M , velocity, & opacity κ



UV/X-ray opacity of neutral oxygen



$$R = v t \quad \rho = \frac{M}{4\pi/3 R^3}$$
$$\tau \sim \kappa \rho R \quad t_{\text{diff}} \sim \tau R/c$$

Light escapes when $t > t_{\text{diff}} \Rightarrow$
 $t > \text{month} \left(\frac{v}{10^4 \text{ km s}^{-1}} \right)^{-1/2} \left(\frac{M}{3M_\odot} \right)^{1/2} \left(\frac{\kappa}{0.1 \text{ cm}^2 \text{g}^{-1}} \right)^{1/2}$


Optical ($\kappa \sim 0.1 \text{ cm}^2 \text{g}^{-1}$) $\Rightarrow t \sim \text{month}$

Hard X-ray $\sim 10 \text{ keV}$ ($\kappa \sim 30 \text{ cm}^2 \text{g}^{-1}$)
 $\Rightarrow t \sim \text{few years}$


Soft X-ray $\sim 0.1 \text{ keV}$ ($\kappa \sim 10^5 \text{ cm}^2 \text{g}^{-1}$)
 $\Rightarrow t \sim 100 \text{ years}$

X-ray Ionization Break-Out

(BDM, Vurm, Hascoet & Beloborodov 2013)



1. Pulsar inflates cavity (pulsar wind nebula)
2. Nebula X-rays ionize inner exposed surface of ejecta
3. Ionization front reaches outer surface - X-rays escape to observer.



Evolution of Millisecond Pulsar Wind Nebulae

(BDM, Vurm, Hascoet & Beloborodov 2013)

Non-Thermal UV / X-rays

Source: cooling $e^{+/-}$ pairs (pulsar)

Sinks: PdV work, absorption by ejecta walls

Thermal Bath (Optical)


Source: re-emission of X-rays by ejecta walls


Sinks: PdV work, radiative diffusion

Ejecta Ionization State

- Balance photo-ionization with recombination in ionized layer(s)
- Sets ejecta albedo (thermalization efficiency)

analogy to AGN accretion disks





Evolution of Millisecond Pulsar Wind Nebulae

(BDM, Vurm, Hascoet & Beloborodov 2013)

Non-Thermal UV / X-rays

Source: cooling $e^{+/-}$ pairs (pulsar)

Sinks: PdV work, absorption by ejecta walls

Thermal Bath (Optical)

Source: re-emission of X-rays by ejecta walls



Sinks: PdV work, radiative diffusion

Ejecta Ionization State





- Balance photo-ionization with recombination in ionized layer(s)
- Sets ejecta albedo (thermalization efficiency)

analogy to AGN accretion disks

Example: $B = 10^{13}$ G, $P = 1$ ms, $M_{ej} = 3 M_{\odot}$





Example: $B = 10^{13}$ G, $P = 1$ ms, $M_{ej} = 3 M_\odot$



Superluminous X-rays from a Superluminous SN

(Levan, Read, BDM, Wheatley, Tanvir 2013)



Chatzopoulos et al. 2013
(cf. Barbary et al. 2009)




Superluminous X-rays from a Superluminous SN

(Levan, Read, BDM, Wheatley, Tanvir 2013)

Chatzopoulos et al. 2013
(cf. Barbary et al. 2009)



Levan et al. 2013



- No detections from other SLSNe
- Upper limits $L_x < 10^{42}-10^{44}$ erg s^{-1} on timescales < 70 days (usually too early!)
- Future: X-ray follow-up *after* optical peak confirm or constrain pulsar model for SLSNe

Summary

- Rapid (millisecond) birth period may be key to generating large scale magnetar-strength B fields.
- Powerful outflow ($\tau_{\text{sd}} \sim \text{min-hour}$) \Rightarrow relativistic jet \Rightarrow GRB
 - Baryon loading set by neutrino heating above magnetar surface.
 - Accounts for GRB energetics, Lorentz factors, duration, collimation, late activity; natural association with energetic supernovae.
 - Key issues: stability of 3D jet and predicted rise in magnetization during GRB.
- Weaker outflow ($\tau_{\text{sd}} \sim \text{weeks}$) \Rightarrow jet trapped \Rightarrow SuperLuminous SN
 - Previous models assume pulsar wind thermalizes with 100% efficiency.
 - We have developed a model for the evolution of young msPWNe that couples X-ray and thermal radiation via interaction with ejecta walls
 - Pulsar wind $\Rightarrow e^{+/-}$ pairs \Rightarrow X-rays \Rightarrow thermal (optical) photons \Rightarrow observer (optical SN)
 - Nebular UV/X-rays can re-ionize ejecta within months of optical peak ('Ionization Break-Out'), allowing escape of high energy radiation.