


# From Raw Data to Physics: Reconstruction and Analysis


Introduction

Sample Analysis

A Model


Basic Features





**We use experiments to inquire about what “reality” does.**

**We intend to fill this gap**



The goal is to understand in the most general; that’s usually also the simplest.  
- A. Eddington

# Theory

146 10. Electroweak model and constraints on new physics

## 10. ELECTROWEAK MODEL AND CONSTRAINTS ON NEW PHYSICS

Revised August 1999 by J. Erler and P. Langacker (Univ. of Pennsylvania).

- 10.1 Introduction
- 10.2 Renormalization and radiative corrections
- 10.3 Cross-section and asymmetry formulas
- 10.4  $W$  and  $Z$  decays
- 10.5 Experimental results
- 10.6 Constraints on new physics

### 10.1. Introduction

The standard electroweak model is based on the gauge group [1]  $SU(2) \times U(1)$ , with gauge bosons  $W_\mu^i$ ,  $i = 1, 2, 3$ , and  $B_\mu$  for the  $SU(2)$  and  $U(1)$  factors, respectively, and the corresponding gauge coupling constants  $g$  and  $g'$ . The left-handed fermion fields  $\psi_i = \begin{pmatrix} \nu_i \\ \ell_i^- \end{pmatrix}$  and  $\begin{pmatrix} u_i \\ d_i \end{pmatrix}$  of the  $i^{\text{th}}$  fermion family transform as doublets under  $SU(2)$ , where  $d_i' \equiv \sum_j V_{ij} d_j$ , and  $V$  is the Cabibbo-Kobayashi-Maskawa mixing matrix. (Constraints on  $V$  are discussed in the section on the Cabibbo-Kobayashi-Maskawa mixing matrix.) The right-handed fields are  $SU(2)$  singlets. In the minimal model there are three fermion families and a single complex Higgs doublet  $\phi \equiv \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ .

After spontaneous symmetry breaking the Lagrangian for the fermion fields is

$$\begin{aligned} \mathcal{L}_F = & \sum_i \bar{\psi}_i \left( i \not{\partial} - m_i - \frac{g m_i H}{2 M_W} \right) \psi_i \\ & - \frac{g}{2\sqrt{2}} \sum_i \bar{\psi}_i \gamma^\mu (1 - \gamma^5) (T^+ W_\mu^+ + T^- W_\mu^-) \psi_i \\ & - e \sum_i q_i \bar{\psi}_i \gamma^\mu \psi_i A_\mu \\ & - \frac{g}{2 \cos \theta_W} \sum_i \bar{\psi}_i \gamma^\mu (g_V^i - g_A^i \gamma^5) \psi_i Z_\mu. \end{aligned} \quad (10.1)$$

$\theta_W \equiv \tan^{-1}(g'/g)$  is the weak angle;  $e = g \sin \theta_W$  is the positron electric charge; and  $A \equiv B \cos \theta_W + W^3 \sin \theta_W$  is the (massless) photon field.  $W^\pm \equiv (W^1 \mp i W^2)/\sqrt{2}$  and  $Z \equiv -B \sin \theta_W + W^3 \cos \theta_W$  are the massive charged and neutral weak boson fields, respectively.  $T^+$  and  $T^-$  are the weak isospin raising and lowering operators. The

Particle Data Group,  
Barnett et al

“Clear statement of how the world works”

Additional term goes here

# Experiment

```
0x01e84c10: 0x01e8 0x8848 0x01e8 0x83d8 0x6c73 0x6f72 0x7400 0x0000
0x01e84c20: 0x0000 0x0019 0x0000 0x0000 0x01e8 0x4d08 0x01e8 0x5b7c
0x01e84c30: 0x01e8 0x87e8 0x01e8 0x8458 0x7061 0x636b 0x6167 0x6500
0x01e84c40: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84c50: 0x01e8 0x8788 0x01e8 0x8498 0x7072 0x6f63 0x0000 0x0000
0x01e84c60: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84c70: 0x01e8 0x8824 0x01e8 0x84d8 0x7265 0x6765 0x7870 0x0000
0x01e84c80: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84c90: 0x01e8 0x8838 0x01e8 0x8518 0x7265 0x6773 0x7562 0x0000
0x01e84ca0: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84cb0: 0x01e8 0x8818 0x01e8 0x8558 0x7265 0x6e61 0x6d65 0x0000
0x01e84cc0: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84cd0: 0x01e8 0x8798 0x01e8 0x8598 0x7265 0x7475 0x726e 0x0000
0x01e84ce0: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84cf0: 0x01e8 0x87ec 0x01e8 0x85d8 0x7363 0x616e 0x0000 0x0000
0x01e84d00: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84d10: 0x01e8 0x87e8 0x01e8 0x8618 0x7365 0x7400 0x0000 0x0000
0x01e84d20: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84d30: 0x01e8 0x87a8 0x01e8 0x8658 0x7370 0x6c69 0x7400 0x0000
0x01e84d40: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84d50: 0x01e8 0x8854 0x01e8 0x8698 0x7374 0x7269 0x6e67 0x0000
0x01e84d60: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84d70: 0x01e8 0x875c 0x01e8 0x86d8 0x7375 0x6273 0x7400 0x0000
0x01e84d80: 0x0000 0x0019 0x0000 0x0000 0x0000 0x0000 0x01e8 0x5b7c
0x01e84d90: 0x01e8 0x87c0 0x01e8 0x8718 0x7377 0x6974 0x6368 0x0000
```

## **1/30th of an event in the BaBar detector**

- Get about 100 events/second

# What does the data mean?

Digitization:



“Address”: what detector element took the reading


“Value”: What the electronics wrote down


Look up type, calibration info

Check valid, convert to useful units/form

Look up/calculate spatial position


Draw





**The imperfect measurement of  
a (set of) interactions in the detector**

**A unique happening:  
Run 21007, event 3916 which  
contains a  $Z \rightarrow \text{xx}$  decay**



**A small number of general equations, with specific  
input parameters (perhaps poorly known)**

# Phenomenology

**A good theory contains very few numbers**

**But it can predict a large number of reactions**

**Getting those predictions from the theory is called “phenomenology”**

## 10.4. *W* and *Z* decays

The partial decay width for gauge bosons to decay into massless fermions  $f_1\bar{f}_2$  is

$$\Gamma(W^+ \rightarrow e^+\nu_e) = \frac{G_F M_W^3}{6\sqrt{2}\pi} \approx 226.5 \pm 0.3 \text{ MeV} \quad , \quad (10.41a)$$

$$\Gamma(W^+ \rightarrow u_i\bar{d}_j) = \frac{CG_F M_W^3}{6\sqrt{2}\pi} |V_{ij}|^2 \approx (707 \pm 1) |V_{ij}|^2 \text{ MeV} \quad , \quad (10.41b)$$

$$\Gamma(Z \rightarrow \psi_i\bar{\psi}_i) = \frac{CG_F M_Z^3}{6\sqrt{2}\pi} [g_V^{i2} + g_A^{i2}] \quad (10.41c)$$


$$\approx \begin{cases} 300.3 \pm 0.2 \text{ MeV} (u\bar{u}), & 167.24 \pm 0.08 \text{ MeV} (\nu\bar{\nu}), \\ 383.1 \pm 0.2 \text{ MeV} (d\bar{d}), & 84.01 \pm 0.05 \text{ MeV} (e^+e^-), \\ 375.9 \mp 0.1 \text{ MeV} (b\bar{b}). \end{cases}$$

From Particle  
Data Book

**Our modified theory predicts a different rate for  $Z \rightarrow \mu\mu$**

- This gives us a way to prove or disprove it!





**The imperfect measurement of  
a (set of) interactions in the detector**

**A unique happening:  
Run 21007, event 3916 which  
contains a  $Z \rightarrow \text{xx}$  decay**

**Specific lifetimes, probabilities, masses,  
branching ratios, interactions, etc**

**A small number of general equations, with specific  
input parameters (perhaps poorly known)**


## A simple analysis: What's BR(Z<sup>0</sup>→μ<sup>+</sup>μ<sup>-</sup>)?


**Measure:**

$$BR(Z^0 \rightarrow \mu^+ \mu^-) = \frac{\text{Number of } \mu^+ \mu^- \text{ events}}{\text{Total number of events}}$$


**Take a sample of events, and count those with a μ<sup>+</sup>μ<sup>-</sup> final state.**

- Two tracks, approximately back-to-back with the expected |p|  
Other kinds of events have more
- Expected energy in calorimeter  
Electrons will deposit most of their energy early in the calorimeter
- Right number of muon hits  
Muons are very penetrating, travel through entire detector







Not  $Z \rightarrow \mu + \mu^-$




Not  $Z \rightarrow \mu + \mu^-$







Not  $Z \rightarrow \mu + \mu^-$




$Z \rightarrow \mu^+ \mu^-$









Not  $Z \rightarrow \mu + \mu^-$




Not  $Z \rightarrow \mu^+ \mu^-$




$Z \rightarrow \mu^+ \mu^-$




Not  $Z \rightarrow \mu + \mu^-$



Not  $Z \rightarrow \mu + \mu^-$





Not  $Z \rightarrow \mu^+ \mu^-$




Not  $Z \rightarrow \mu^+ \mu^-$







Not  $Z \rightarrow \mu^+ \mu^-$



Not  $Z \rightarrow \mu + \mu^-$

## Summary so far

We have a result:  $BR(Z \rightarrow \mu^+ \mu^-) = 2/45$

But there's a lot more to do!

### Statistical error

- We saw 2 events, but it could easily have been 1 or 3
- Those fluctuations go like the square-root of the number of events:

$$BR(Z^0 \rightarrow \mu^+ \mu^-) = \frac{N_{\mu\mu}}{N_{total}} \pm \frac{\sqrt{N_{\mu\mu}}}{N_{total}}$$

- To reduce that uncertainty, you need lots of events  
Need to record lots of events in the detector, and then process them


### Systematic error

- What if you only see 50% of the  $\mu^+ \mu^-$  events?  
Due to detector imperfections, poor understanding, etc?

$$N_{\mu\mu \text{ seen}} = \epsilon N_{\mu\mu}$$

$$BR(Z^0 \rightarrow \mu^+ \mu^-) = \frac{N_{\text{seen}} / \epsilon}{N_{total}}$$

## Our model so far...



We “confront theory with experiment” by comparing what we measured, with what we expected from our hypothesis.

## The process in practice:


**The reconstruction step is usually done in common**

- “Tracks”, “particle ID”, etc are general concepts, not analysis-specific. Common algorithms make it easier to understand how well they work.
- Common processing needed to handle large amounts of data. Data arrives every day, and the processing has to keep up.

**Analysis is a very individual thing**

- Many different measurements being done at once
- Small groups working on topics they’re interested in
- Many different timescales for these efforts


**Collaborations build “offline computing systems” to handle all this.**



# Reconstruction: Calorimeter Energy

**Goal is to measure particle properties in the event**

- “Finding” stage attempts to find patterns that indicate what happened
- “Fitting” stage attempts to extract the best possible measurement from those patterns.




## Finding

**Clusters of energy in a calorimeter are due to the original particles**

- Clustering algorithm groups individual channel energies
- Don't want to miss any; don't want to pick up fakes

**Many algorithms exist**


- Scan for one or more channels above a high threshold as “seeds”
- Include channels on each side above a lower threshold:



**Not perfect! Doesn't use prior knowledge about event, cluster shape, etc**



# One lump or two?



**Hard to tune thresholds to get this right.**

**Perhaps a smarter algorithm would do better?**

# Fitting

From the clusters, fit for energy and position

- Complicated by noise & limited information

Simple algorithm: Sum of channels for energy, average for position





Figure 8 Correlation between the positions measured with (a) the center of gravity method ( $X_{cg}$ ) and (b) the reconstructed positions ( $X_{rec}$ ) vs the actual positions ( $X_{in}$ ). The results are derived from 5000  $Z \rightarrow e^+e^-$  decays simulated by the GEANT Monte Carlo in the L3 HGO calorimeter (44).

**Empirical corrections are important!**

## Analysis: Measure BR(B $\rightarrow$ J/ $\Psi$ K $\rightarrow$ K $\pi$ )

Neither J/ $\Psi$  nor K $\rightarrow$ K $\pi$  is a long-lived particle

- Detector doesn't see them, only their decay products K $\rightarrow$ K $\pi$


Take all pairs of possible particles, and calculate their mass

$$m^2 = E^2 - p^2 = (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2$$


If it's not the K $\rightarrow$ K $\pi$  mass,  
that combination can't be a K $\rightarrow$ K $\pi$

If it is the K $\rightarrow$ K $\pi$  mass, it  
might be a K $\rightarrow$ K $\pi$


Signal/Background ratio  
is critical to success!



## Next, look for $J/\psi \rightarrow e^+e^-$ and $J/\psi \rightarrow \mu^+\mu^-$



$M_{J/\psi} \rightarrow ee$



$M_{J/\psi} \rightarrow \mu\mu$

### **Why not $J/\psi \rightarrow$ hadrons? Too many wrong combinations!**


- Only a few e/m in an event, so only a few combinations
- About 10 hadrons, so about 50 combinations of two

Some are bound to be at about the right mass!

### **Note peaks not same size, shape**

- Do we understand our efficiency?

# Monte Carlo simulation's role



# How do you know it is correct?

## Divide and conquer

- A very detailed simulation can reproduce even unlikely problems
- By making it of small parts, each can be understood
- Some aspects are quite general, so detailed handling is possible

## Why does it matter?

- We “cut on” distributions
- Example: Energy (e.g. signal) from particle in a Si detector




Fig. 15 Comparison of measured and simulated energy deposition in 530  $\mu\text{m}$  silicon for 1 MeV electrons (experimental points see [30]).

Take only particles to left of blue line

Dots are data in test beam

Two solid lines are two simulation codes

One simulation doesn't provide the right efficiency!






Figure 18: Observed mass distribution superimposed with uds, cc, generic  $B\bar{B}$  and signal MC events for (a)  $J/\psi \rightarrow e^+e^-$  and (b)  $J/\psi \rightarrow \mu^+\mu^-$ .

**The tricky part is understanding the discrepancies....**

# Finally, put together parts to look for $B \rightarrow J/\psi K^*$

## Details:

- Background under peak?
- Systematic errors on efficiency
- .....



**When you get more data, you need to do a better job on the details**



## Summary so far

We seen some simple analyses

We have a model of the steps involved

We're starting to see details of how its done

More detailed examples tomorrow!

