

# Physics 56400

# Introduction to Elementary Particle Physics I

*Now in PowerPoint!*

Lecture 2  
Fall 2018 Semester  
Prof. Matthew Jones

# Cross Sections

- Reaction rate:

$$R = \mathcal{L} \cdot \sigma$$

- The cross section is proportional to the probability of observing a particular outcome in a collision
- Example:
  - High energy protons might interact with a large cross section but only rarely do they produce a Higgs particle
  - The rates of both processes are proportional to the luminosity

# Cross Sections

- In quantum mechanics, one of the only questions we are allowed to ask is,

“what is the probability of observing a system in a final state  $|f\rangle$  given that it started in an initial state  $|i\rangle$ ?”

- Quantum mechanics states that this probability is proportional to

$$P \sim |\langle f|i\rangle|^2$$

- If the initial and final states describe “free particles”, then

$$P \sim |\langle f|U|i\rangle|^2$$

where the operator  $U$  evolves the initial state, from time  $t \rightarrow -\infty$  to the asymptotic time  $t \rightarrow +\infty$ .

# Cross Sections


- We will see that cross sections work like probabilities:

$$\sigma \sim |\langle f|U|i\rangle|^2$$

- The time evolution operator is expressed in terms of the Hamiltonian,  $H$ .
- We are now allowed to ask *how* the initial state turns into the final state, we can only calculate the probability.
- In practice, we need to account for all possible intermediate states, and add their amplitudes (complex numbers).
- The Standard Model of particle physics tells us what the possible intermediate states could be and how the initial and final states couple to them.


# Cross Sections

- The cross section could be a function of several independent variables
  - For example, the beam energy:



# Cross Sections

The deuteron is a bound state of a proton and a neutron.  
It is a stable isotope of hydrogen (heavy hydrogen).




# Differential Cross Sections

- There are lots of ways to describe the final state
- The total proton-proton cross section accounts for all possible interactions
- The elastic cross section describes the process where the protons retain their identity but just “bounce” off of each other
- We detect these by observing protons scattered at some angle with respect to the initial beam direction
- What can we learn by measuring the angular distribution of scattered protons?

# Geometry

- We will use spherical coordinates to describe the scattering geometry:



- The angle  $\theta$  is measured with respect to the  $z$ -axis
  - The angle  $\varphi$  is measured with respect to the  $x$ -axis in the  $x$ - $y$  plane.
- (This is just the usual system of polar coordinates)
- In systems with azimuthal symmetry, we don't expect there to be any dependence on  $\varphi$ .



# Geometry

- The differential element of solid angle is

$$d\Omega = d(\cos \theta) d\varphi = \sin \theta d\theta d\varphi$$

- This solid angle has the area  $dA$  on the surface of a sphere of radius  $R$ :

$$dA = R^2 d\Omega$$

- The total surface area of the sphere is

$$\begin{aligned} A &= R^2 \int_{\pi}^0 \sin \theta d\theta \int_0^{2\pi} d\varphi = 2\pi R^2 \int_{-1}^1 d(\cos \theta) \\ &= 4\pi R^2 \end{aligned}$$

# Differential Cross Sections


- We want to measure the rate at which protons are scattered with polar angles  $(\theta, \varphi)$  into a small interval of solid angle:

$$dR = \mathcal{L} \frac{d\sigma}{d\Omega} d\Omega$$

- The function  $d\sigma/d\Omega$  is the differential cross section.
  - In this case it is a function of the polar angles
- Let's calculate  $d\sigma/d\Omega$  for a couple different models...

# Brick Wall Scattering


- Suppose the target particles were flat disks:



- Assuming they were all oriented the same way, the target would just reflect the beam particles
- All scattering angles would be the same
- The differential cross section would be a delta-function at that specific scattering angle  
(not very interesting or realistic)

# Hard Sphere Scattering

- Given that we are probably scattering from nuclei, perhaps they could be described as hard spheres...



"Impact parameter",  $b$

$$\sin \alpha = b/R$$

# Hard Sphere Scattering

- The scattering angle is  $\theta = \pi - 2\alpha$

$$\cos \theta = \cos(\pi - 2\alpha)$$

$$= -\cos 2\alpha$$

$$= \sin^2 \alpha - \cos^2 \alpha$$

$$= 2 \sin^2 \alpha - 1$$

$$= \frac{2b^2}{R^2} - 1$$

- Does this make sense?

- As  $b \rightarrow R$ ,  $\cos \theta \rightarrow 1$

- As  $b \rightarrow 0$ ,  $\cos \theta \rightarrow -1$

# Hard Sphere Scattering

- The area of the hard sphere that will result in scattering angles larger than  $\theta$  will be

$$\begin{aligned} A &= \pi b^2 \\ &= \frac{\pi}{2} R^2 (1 + \cos \theta) \end{aligned}$$

- The differential scattering cross section is


$$\frac{d\sigma}{d\Omega} = \frac{d\sigma}{d(\cos \theta) d\varphi} = \frac{R^2}{4}$$

– This is isotropic

- Total cross section is  $\sigma = \pi R^2$

# Coulomb Scattering

- Assume that the beam has charge  $ze$  and the target nuclei has charge  $Ze$
- The beam particle will follow a hyperbolic trajectory:



- Classical mechanics:

$$b = \frac{zZe^2}{mv_0} \cot(\theta/2)$$

But we don't know  $b$  on an event-by-event basis...

# Coulomb Scattering

- Assume that the beam has intensity  $I$ .
- $I$  is the number of incident beam particles per unit area per unit time.
- Total number of beam particles:

$$N = \int_0^T dt \int_0^\infty 2\pi b db I(b, t)$$

- We typically assume that the intensity is independent of time
- By definition,

$$d\sigma = \frac{dN(\theta, \varphi)}{T \cdot I_0} = b db d\varphi$$



# Coulomb Scattering

- Differential cross section:

$$\frac{d\sigma}{d\Omega} = \frac{b \, db \, d\varphi}{d(\cos \theta) \, d\varphi} = \frac{b}{\sin \theta} \frac{db}{d\theta}$$

- But,

$$\begin{aligned} \frac{db}{d\theta} &= \frac{d}{d\theta} \left( \frac{zZe^2}{mv_0} \cot(\theta/2) \right) \\ &= \frac{zZe^2}{mv_0} \csc^2(\theta/2) \end{aligned}$$


# Coulomb Scattering

- Eventually, it turns out that

$$\frac{d\sigma}{d\Omega} = \frac{1}{4} \left( \frac{zZe^2}{mv_0} \right)^2 \frac{1}{\sin^4(\theta/2)}$$

- This is the cross section for scattering from a single target nucleus

$$\frac{dR}{d\Omega} = \mathcal{L} \frac{d\sigma}{d\Omega}$$



# Differential Cross Sections

- By studying the dependence of the differential cross section on various independent variables we can learn about the fundamental interactions on the microscopic scale.
- Exclusive cross section:
  - Final state is specified precisely (eg, exactly one proton in the case of elastic scattering)
- Inclusive cross section:
  - Final state includes the specified particle configuration
  - This is usually the sum of all possible exclusive cross sections that include the specified particle configuration

# Decay Rates

- What else can we measure?
  - Lifetimes of unstable particles
  - Branching fractions to different final states
- Lifetime, decay rate and “partial width”:

$$\Gamma = \frac{\hbar}{\tau}$$

- Branching fraction:

$$Br(i \rightarrow f) = \frac{\Gamma_f}{\Gamma_{total}}$$

$$\Gamma_{total} = \sum_j \Gamma_j$$

$$\Gamma_j \sim |\langle j|U|i\rangle|^2$$