

Multiple choice (5 pts each) – circle the correct answers.

1. Sagittarius A* is a black hole with mass 4.3×10^6 solar masses while Messier 108 is a black hole with mass 2.4×10^7 solar masses. What is the diameter of Sagittarius A* relative to Messier 108?

(a) $5.6 \times$ larger	(b) $5.6 \times$ smaller	(c) $31 \times$ larger
(d) $31 \times$ smaller	(e) $174 \times$ larger	(f) $174 \times$ smaller

2. Light from a laser with wavelength λ and intensity $1 \mu\text{J}/\text{m}^2$ has 3.2×10^8 photons per second go through an area of 1 cm^2 . For a laser with the same intensity but wavelength $\lambda/10$, how many photons go through an area of 1 cm^2 ?

(a) 3.2×10^7 photons per second
(b) 3.2×10^8 photons per second
(c) 3.2×10^9 photons per second

3. You have found two solutions of Schrodinger's equation, $\Psi_1(x, t)$ and $\Psi_2(x, t)$. Which one is also a possible solution?

(a) $\Psi(x, t) = C_1\Psi_1(x, t) + C_2\Psi_2(x, t)$ where C_1 and C_2 don't depend on x or t
(b) $\Psi(x, t) = C_1(x)\Psi_1(x, t) + C_2(x)\Psi_2(x, t)$ where C_1 and C_2 must depend on x
(c) $\Psi(x, t) = C_1(t)\Psi_1(x, t) + C_2(t)\Psi_2(x, t)$ where C_1 and C_2 must depend on t
(d) $\Psi(x, t) = C_1(x, t)\Psi_1(x, t) + C_2(x, t)\Psi_2(x, t)$ where C_1 and C_2 must depend on x and t

4. A beam of photons with an energy of 6.78 eV is incident on aluminum and absorbed. Will electrons be ejected? If yes, give their maximum kinetic energy.

(a) no	(b) yes, 4.50 eV	(c) yes, 2.70 eV
(d) yes, 2.08 eV	(e) yes, 6.78 eV	(f) yes, 10.86 eV

5. Astronauts leave earth and travel to a star 4 light-years away and immediately come back. The space ship they were in traveled at $0.8 c$. How much did the astronauts age?

(a) less than 10 years
(b) 10 years
(c) more than 10 years

6. You measure that 2 galaxies are moving away from the Milky Way. The first moves away with speed 10^7 m/s and the second moves away with speed $2 \times 10^7 \text{ m/s}$. The distance to the second galaxy is

(a) roughly 1/2 the distance to the first.
(b) roughly the same distance to the first.
(c) roughly 2 times the distance to the first.

Problems: Show *all* work to receive full credit.

Fundamental constants (e , h , masses, etc) are top of the equation page.

- 10 pts 1. A star with radius 9.0×10^8 m radiates light like an ideal thermal source at a temperature of 7200 K. a) Keeping the temperature fixed, how would the rate that energy is radiated change if the radius changes to 7.7×10^8 m? b) Keeping the radius fixed, how would the rate that energy is radiated change if the temperature increases to 8100 K?

- 10 pts 2. A spaceship is traveling directly at you at a speed of $0.500\ c$. They fire at you a proton beam. You measure the protons in the beam to have a speed of $0.950\ c$. a) What is the speed of the protons as measured by the creatures on the spaceship? b) In firing one proton, what is the *change* in momentum of the spaceship as measured by you?

10 pts 3. A type of Kaon with mass $497.611 \text{ MeV}/c^2$ decays into two identical Pions (the mass of one Pion is $134.977 \text{ MeV}/c^2$). If the Kaon is initially at rest, what speed does each Pion have?

10 pts 4. An electron travels in a region with no forces. It has speed of $3.0 \times 10^6 \text{ m/s}$. Determine the electron's wavelength and frequency.

- 15 pts 5. Light with a wavelength of 633 nm and intensity of 1.30 W/m^2 is incident on a spherical microparticle. Assume the light travels in the $+y$ -direction and the microparticle is in space with no other interactions. The radius of the microparticle is $4.5 \times 10^{-6} \text{ m}$ and it has a density of 1800 kg/m^3 . Assume that all of the photons that hit the microparticle are absorbed. a) Does the mass of the microparticle increase with time? If so, what is the rate of change of the mass (give your answer in kg/s). b) Does the microparticle accelerate? If so, give the acceleration vector.

- 15 pts 6. An electron experiences a potential energy which is $U(x) = 0$ for $x_0 < x < x_f$ and $U(x) = \infty$ for $x < x_0$ or $x > x_f$. The x_0, x_f are constants. a) What are the allowed wavelengths for the electron? b) What are the allowed momenta? c) If you **don't** have to worry about relativity, what are the allowed energies? d) If you **do** have to worry about relativity, what are the allowed energies? e) For the nonrelativistic case, the electron starts in the 5th state then emits a photon and ends in the 4th state. What is the frequency of the photon?

$$\begin{aligned}
g &= 9.80 \frac{m}{s^2} & h &= 6.63 \times 10^{-34} \text{ J s} = 4.14 \times 10^{-15} \text{ eV s} & \hbar &= \frac{h}{2\pi} & c &= 3.00 \times 10^8 \frac{m}{s} \\
M_{elec} &= 9.11 \times 10^{-31} \text{ kg} & M_{prot} &= 1.67 \times 10^{-27} \text{ kg} & M_{muon} &= 1.88 \times 10^{-28} \text{ kg} \\
1.60 \times 10^{-19} \text{ J} &= 1 \text{ eV} & \text{Work functions: Na } 2.28 \text{ eV, Al } 4.08 \text{ eV, Cu } 4.70 \text{ eV} \\
V_{sph} &= \frac{4}{3}\pi r^3
\end{aligned}$$

Galilean relativity	$x' = x - ut, v'_x = v_x - u$	2.1	Lorentz velocity transformation	$v'_x = \frac{v_x - u}{1 - v_x u / c^2},$	2.5
Einstein's postulates	(1) The laws of physics are the same in all inertial frames. (2) The speed of light has the same value c in all inertial frames.	2.3		$v'_y = \frac{v_y \sqrt{1 - u^2/c^2}}{1 - v_x u / c^2},$	
				$v'_z = \frac{v_z \sqrt{1 - u^2/c^2}}{1 - v_x u / c^2}$	
Time dilation	$\Delta t = \frac{\Delta t_0}{\sqrt{1 - u^2/c^2}}$ (Δt_0 = proper time)	2.4	Clock synchronization	$\Delta t' = \frac{uL/c^2}{\sqrt{1 - u^2/c^2}}$	2.5
Length contraction	$L = L_0 \sqrt{1 - u^2/c^2}$ (L_0 = proper length)	2.4	Relativistic momentum	$\vec{p} = \frac{m\vec{v}}{\sqrt{1 - v^2/c^2}}$	2.7
Velocity addition	$v = \frac{v' + u}{1 + v' u / c^2}$	2.4	Relativistic kinetic energy	$K = \frac{mc^2}{\sqrt{1 - v^2/c^2}} - mc^2$	2.7
Doppler effect (source and observer separating)	$f' = f \sqrt{\frac{1 - u/c}{1 + u/c}}$	2.4	Rest energy	$E_0 = mc^2$	2.7
Lorentz transformation	$x' = \frac{x - ut}{\sqrt{1 - u^2/c^2}},$ $y' = y, z' = z,$ $t' = \frac{t - (u/c^2)x}{\sqrt{1 - u^2/c^2}}$	2.5	Relativistic total energy	$E = K + E_0 = \frac{mc^2}{\sqrt{1 - v^2/c^2}}$	2.7
			Momentum-energy relationship	$E = \sqrt{(pc)^2 + (mc^2)^2}$	2.7
			Extreme relativistic approximation	$E \cong pc$	2.7
			Conservation laws	In an isolated system of particles, the total momentum and the relativistic total energy remain constant.	2.8
Hubble's law	$v = H_0 d$	15.1	Age of matter-dominated universe	$t = 1/\sqrt{6\pi G \rho_m}$	15.7
Number density of photons	$N/V = (2.03 \times 10^7 \text{ photons/m}^3 \cdot \text{K}^3) T^3$	15.2	Age of radiation-dominated universe	$t = \sqrt{3/32\pi G \rho_r}$	15.7
Energy density of photons	$U = (4.72 \times 10^3 \text{ eV/m}^3 \cdot \text{K}^4) T^4$	15.2	Temperature of universe at age t	$T = \frac{1.5 \times 10^{10} \text{ s}^{1/2} \cdot \text{K}}{t^{1/2}}$	15.8
Gravitational frequency change	$\Delta f/f = gH/c^2$	15.4	Fraction of photons above E_0	$f = 0.42 e^{-E_0/kT} \times \left[\left(\frac{E_0}{kT} \right)^2 + 2 \left(\frac{E_0}{kT} \right) + 2 \right]$	15.9
Deflection of starlight	$\theta = 2GM/Rc^2$	15.5	Critical density of universe	$\rho_{cr} = \frac{3H^2}{8\pi G} = 0.97 \times 10^{-26} \text{ kg/m}^3$	15.10
Perihelion precession	$\Delta\phi = \frac{6\pi GM}{c^2 r_{\min}(1 + e)}$	15.5			
Schwarzschild radius	$r_s = 2GM/c^2$	15.6			

Double-slit maxima	$y_n = n \frac{\lambda D}{d} \quad n = 0, 1, 2, 3, \dots$	3.1	Rayleigh-Jeans formula	$I(\lambda) = \frac{2\pi c}{\lambda^4} kT$	3.3
Bragg's law for X-ray diffraction	$2d \sin \theta = n\lambda \quad n = 1, 2, 3, \dots$	3.1	Planck's blackbody distribution	$I(\lambda) = \frac{2\pi hc^2}{\lambda^5} \frac{1}{e^{hc/\lambda kT} - 1}$	3.3
Energy of photon	$E = hf = hc/\lambda$	3.2	Compton scattering	$\frac{1}{E'} - \frac{1}{E} = \frac{1}{m_e c^2} (1 - \cos \theta),$ $\lambda' - \lambda = \frac{h}{m_e c} (1 - \cos \theta)$	3.4
Maximum kinetic energy of photoelectrons	$K_{\max} = eV_s = hf - \phi$	3.2	Bremsstrahlung	$\lambda_{\min} = hc/K = hc/e\Delta V$	3.5
Cutoff wavelength	$\lambda_c = hc/\phi$	3.2	Pair production	$hf = E_+ + E_- = (m_e c^2 + K_+) + (m_e c^2 + K_-)$	3.5
Stefan's law	$I = \sigma T^4$	3.3	Electron-positron annihilation	$(m_e c^2 + K_+) + (m_e c^2 + K_-) = E_1 + E_2$	3.5
Wien's displacement law	$\lambda_{\max} T = 2.8978 \times 10^{-3} \text{ m} \cdot \text{K}$	3.3			

De Broglie wavelength	$\lambda = h/p$	4.1	Statistical momentum uncertainty	$\Delta p_x = \sqrt{(p_x^2)_{\text{av}} - (p_{x,\text{av}})^2}$	4.4
Single slit diffraction	$a \sin \theta = n\lambda \quad n = 1, 2, 3, \dots$	4.2	Wave packet (discrete k)	$y(x) = \sum A_i \cos k_i x$	4.5
Classical position-wavelength uncertainty	$\Delta x \Delta \lambda \sim \varepsilon \lambda^2$	4.3	Wave packet (continuous k)	$y(x) = \int A(k) \cos kx \, dk$	4.5
Classical frequency-time uncertainty	$\Delta f \Delta t \sim \varepsilon$	4.3	Group speed of wave packet	$v_{\text{group}} = \frac{d\omega}{dk}$	4.6
Heisenberg position-momentum uncertainty	$\Delta x \Delta p_x \sim \hbar$	4.4			
Heisenberg energy-time uncertainty	$\Delta E \Delta t \sim \hbar$	4.4			

Time-independent Schrödinger equation	$-\frac{\hbar^2}{2m} \frac{d^2 \psi}{dx^2} + U(x) \psi(x) = E \psi(x)$	5.3	Infinite potential energy well	$\psi_n(x) = \sqrt{\frac{2}{L}} \sin \frac{n\pi x}{L},$ $E_n = \frac{\hbar^2 n^2}{8mL^2} \quad (n = 1, 2, 3, \dots)$	5.4
Time-dependent Schrödinger equation	$\Psi(x, t) = \psi(x) e^{-i\omega t}$	5.3	Two-dimensional infinite well	$\psi(x, y) = \frac{2}{L} \sin \frac{n_x \pi x}{L} \sin \frac{n_y \pi y}{L}$ $E = \frac{\hbar^2}{8mL^2} (n_x^2 + n_y^2)$	5.4
Probability density	$P(x) = \psi(x) ^2$	5.3	Simple harmonic oscillator ground state	$\psi(x) = (m\omega_0/\hbar\pi)^{1/4} e^{-(\sqrt{km}/2\hbar)x^2}$	5.5
Normalization condition	$\int_{-\infty}^{+\infty} \psi(x) ^2 dx = 1$	5.3	Simple harmonic oscillator energies	$E_n = (n + \frac{1}{2}) \hbar \omega_0 \quad (n = 0, 1, 2, \dots)$	5.5
Probability in interval x_1 to x_2	$P(x_1 : x_2) = \int_{x_1}^{x_2} \psi(x) ^2 dx$	5.3	Potential energy step, $E > U_0$	$\psi_0(x < 0) = A \sin k_0 x + B \cos k_0 x$ $\psi_1(x > 0) = C \sin k_1 x + D \cos k_1 x$	5.6
Average or expectation value of $f(x)$	$[f(x)]_{\text{av}} = \int_{-\infty}^{+\infty} \psi(x) ^2 f(x) dx$	5.3	Potential energy step, $E < U_0$	$\psi_0(x < 0) = A \sin k_0 x + B \cos k_0 x$ $\psi_1(x > 0) = C e^{k_1 x} + D e^{-k_1 x}$	5.6
Constant potential energy, $E > U_0$	$\psi(x) = A \sin kx + B \cos kx,$ $k = \sqrt{2m(E - U_0)}/\hbar$	5.4			
Constant potential energy, $E < U_0$	$\psi(x) = A e^{k'x} + B e^{-k'x},$ $k' = \sqrt{2m(U_0 - E)}/\hbar$	5.4			