

Decoherence of rotational degrees of freedom

Changchun Zhong^{1,*} and F. Robicheaux^{1,2,†}¹*Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA*²*Purdue Quantum Center, Purdue University, West Lafayette, Indiana 47907, USA*

(Received 9 May 2016; published 11 November 2016)

The mechanism of decoherence for a mesoscopic quantum system with rotational degrees of freedom is studied. From a simple model of elastic scattering, we show that the nondiagonal density-matrix elements of the system exponentially decay. The decay rate depends on the difference of scattering amplitudes for different rotational configurations, leading to the gradual loss of quantum coherence between the pointer states in the orientational space. For a dielectric ellipsoid immersed in a photon-gas environment (assuming no absorption), the decay rate is found to be proportional to the seventh power of the temperature. For an ellipsoidal object interacting with massive particles, the decay rate is proportional to the 5/2 power of the temperature. Both are different from the case of translational decoherence induced by the same environment scattering. For photon scattering, the coherence time in the rotational degrees of freedom is shown to be much shorter than that in the translational degrees of freedom.

DOI: [10.1103/PhysRevA.94.052109](https://doi.org/10.1103/PhysRevA.94.052109)

I. INTRODUCTION

Decoherence refers to the process of a quantum system losing its quantum coherence between pointer states. Decoherence is not only of great importance to the foundations of quantum physics [1–4], but also of vital interest for the realization of quantum applications, such as quantum computers [5], and other coherent manipulations [6], etc. The key idea of decoherence is simple and clear, that a quantum system in reality is essentially open because it could never be completely isolated from its environment. Thus, this open quantum system is not expected to follow the Schrödinger equation, instead, it evolves nonunitarily according to a master equation [1,2]. For large quantum systems, the density matrix will quickly become mixed, causing the quantum to classical transition [3,4,7].

With fast technological developments in all areas of physics, quantum phenomena become controllable at larger and larger scales [8–13]. For large systems, the rotational or vibrational degrees of freedom is significant. However, in the past decades, most of the attention of mesoscopic systems has been concentrated on decoherence of the translational degrees of freedom [2,14,15], where environmental interactions can produce exponentially localized wave packets. In this paper, we focus on the mechanism of decoherence for a mesoscopic system with rotational degrees of freedom. The answer to these questions should provide guidance in the growing number of experiments probing and controlling the internal degrees of freedom of a mesoscopic system [16].

In a recent work [17], Fischer considered the interaction of a rigid body with an environment through the coupling of a dipole moment and a fluctuating field. Based on the method of quantum stochastic differential equation (SDE), Fischer derived a master equation

$$\partial_t \rho \propto \int_0^\infty ds \iint d\hat{n}_1 d\hat{n}_2 \gamma(s) (e^{i f_s(\hat{\Omega})} \rho e^{-i f_s(\hat{\Omega})} - \rho), \quad (1)$$

which describes the dynamics of an extended object interacting with random pulsed fields (the unitary part is not included). ρ is the system density operator, s is random number, \hat{n}_1, \hat{n}_2 give the direction of the dipole (in the body-fixed frame) and the field (in the space-fixed frame of reference), $\gamma(s)$ is proportional to a normalized distribution function, and $e^{i f_s(\hat{\Omega})}$ is the jump operator. The master equation gives an exponential decay of angular coherence, with the decay rate proportional to the distance measure in orientational space. In Ref. [17], it is worth mentioning that the master equation was also used to discuss the pointer states of the orientational decoherence, and solitonic solutions were identified as the pointer states, which is instrumental for the analysis of rotational dynamics.

In this paper, we extend the first results of Ref. [17] to include rotational decoherence due to interactions with an environment of discrete particles. The derivation is based on a model of scattering which has been widely applied to the decoherence for translational degrees of freedom [1,2,14,15,18]. The single scattering event is treated in a nonperturbative way [2,14,15]. If the system is initially prepared in a rotational superposition state, the coherence in the density matrix is effectively decreased by scattering. For a spherically symmetric environment, the rotational decoherence rate is found to depend only on the difference of the scattering amplitudes for different rotational configurations and can be shown to only depend on the difference in the angles of orientation. To illustrate the effect, we calculate the decoherence rate for the situation with the environment being a photon gas or massive particles. In the long-wavelength limit, the rotational decoherence rate is found to have a temperature dependence different from that for translational decoherence. For photon scattering by systems of the same size, the rotational decoherence rate is shown to be much larger than that in the translational degrees of freedom.

In the sections that follow, we first introduce the derivation of rotational decoherence, present the examples of a system immersed in a photon gas or massive particle environment, compare the result to our expectations, and comment on their implications and possible guidance for future experiments.

*zchangch@purdue.edu

†robichf@purdue.edu

II. ROTATIONAL DECOHERENCE DUE TO SCATTERING

We focus on the rotational degrees of freedom of an object (system S), which interacts with the environment (environment E), and they are assumed to be initially uncorrelated. The combined system (SE) is described by a product state

$$\hat{\rho}_{SE} = \hat{\rho}_S \otimes \hat{\rho}_E, \quad (2)$$

where $\rho_{(S)}$ is the system density operator. ρ_E describes the environment state, which is discussed below. We denote the orientational eigenstate of the system by $|\Omega\rangle = |\alpha, \beta, \gamma\rangle$ [17,19], where α , β , and γ are the Euler angles. In the orientation space, the system density operator takes the form

$$\hat{\rho}_S = \int d\Omega \int d\Omega' \rho_S(\Omega, \Omega') |\Omega\rangle \langle \Omega'|. \quad (3)$$

In the following, we will show how the system state is affected by a single scattering event. The derivation formally follows the steps of the model of collisional decoherence [1,2,20].

A. Correlation established by scattering

Our discussion is confined to elastic scattering throughout the paper. The state $|\Omega\rangle = |\alpha, \beta, \gamma\rangle$ can be represented as a state $|0,0,0\rangle$ rotated by an operator

$$|\Omega\rangle = \hat{D}_S(\Omega)|0,0,0\rangle, \quad (4)$$

where $\hat{D}_S(\Omega) = \exp(-\frac{i}{\hbar}\hat{L}_z\alpha)\exp(-\frac{i}{\hbar}\hat{L}_y\beta)\exp(-\frac{i}{\hbar}\hat{L}_z\gamma)$ [21]. If we denote the incoming particle by $|\chi\rangle$, then the effect of the scattering event can be described by the scattering operator \hat{S} acting on the initial state,

$$|\Omega\rangle |\chi\rangle \rightarrow \hat{S} |\Omega\rangle |\chi\rangle. \quad (5)$$

Then we have

$$\begin{aligned} \hat{S} |\Omega\rangle |\chi\rangle &\rightarrow \hat{S} \hat{D}_{SE}(\Omega) |0,0,0\rangle \hat{D}_E^\dagger(\Omega) |\chi\rangle \\ &\rightarrow \hat{D}_{SE}(\Omega) \hat{S} |0,0,0\rangle \hat{D}_E^\dagger(\Omega) |\chi\rangle, \end{aligned} \quad (6)$$

where $\hat{D}_E(\Omega)$ is the rotational operator acting only on the environment and $\hat{D}_{SE}(\Omega)$ is the rotational operator for the combined system. Obviously, we have $\hat{D}_{SE}(\Omega) = \hat{D}_S(\Omega) \hat{D}_E(\Omega)$. In the last line, we used the fact that the scattering operator commutes with the rotation of the combined system

$$[\hat{S}, \hat{D}_{SE}(\Omega)] = 0. \quad (7)$$

In the scattering model, an important step is to include the nonrecoil approximation, which states that the scattering event essentially does not disturb the system, except establishing entanglement between the system and the incoming particle [2]. In the situations we are considering, the system is much more massive than the environment particles, such as photons scattered by a mesoscopic dielectric or air molecules scattered by more massive objects. This justifies the nonrecoil approximation, which gives

$$\begin{aligned} \hat{S} |\Omega\rangle |\chi\rangle &\rightarrow \hat{D}_{SE}(\Omega) |0,0,0\rangle \hat{S} \hat{D}_E^\dagger(\Omega) |\chi\rangle \\ &\rightarrow |\Omega\rangle \hat{S}(\Omega) |\chi\rangle \rightarrow |\Omega\rangle |\chi(\Omega)\rangle, \end{aligned} \quad (8)$$

where $\hat{S}(\Omega) = \hat{D}_E(\Omega) \hat{S} \hat{D}_E^\dagger(\Omega)$. $|\chi(\Omega)\rangle = \hat{S}(\Omega) |\chi\rangle$ is introduced to denote the state of the outgoing particle, which now carries the orientational information of the system. The first

line used the nonrecoil approximation. From Eq. (8), we see the scattering event establishes correlations between the system and the environment. Accordingly, the initial separable density matrix of the combined system $\rho_{SE} = \rho_S \otimes \rho_E$ is transformed into the following entangled density matrix,

$$\rho_{SE} = \int d\Omega \int d\Omega' \rho_S(\Omega, \Omega') |\Omega\rangle \langle \Omega'| \otimes |\chi(\Omega)\rangle \langle \chi(\Omega')|, \quad (9)$$

where $\rho_S(\Omega, \Omega')$ is typically nondiagonal. The system is described by a reduced density matrix ρ_S , which is obtained by tracing over the environmental degree of freedom,

$$\text{tr}_E(\rho_{SE}) = \int d\Omega \int d\Omega' \rho_S(\Omega, \Omega') |\Omega\rangle \langle \Omega'| \langle \chi(\Omega') | \chi(\Omega)\rangle. \quad (10)$$

As a result, the density-matrix element of the system after the scattering event becomes

$$\rho_S(\Omega, \Omega', 0) \rightarrow \rho_S(\Omega, \Omega', 0) \langle \chi(\Omega') | \chi(\Omega)\rangle, \quad (11)$$

where $\langle \chi(\Omega') | \chi(\Omega)\rangle = \langle \chi | \hat{S}^\dagger(\Omega') \hat{S}(\Omega) | \chi \rangle$. Thus, a suppression is attached to the system density-matrix elements, and the value is determined by the average of the operator $\hat{S}^\dagger(\Omega') \hat{S}(\Omega)$ over the state of the incoming particle. The overlap $\langle \chi(\Omega') | \chi(\Omega)\rangle$ is trivially one for $\Omega = \Omega'$, which indicates no influence on the diagonal elements from the scattering.

B. Time evolution of the system density matrix

To derive how the system density matrix evolves in time, we first need to calculate the overlap $\langle \chi(\Omega') | \chi(\Omega)\rangle$ and then determine how the system density matrix is affected by successive scattering events. According to Eq. (11), the suppression of the system density-matrix element is determined by the average of the operator $\hat{S}^\dagger(\Omega') \hat{S}(\Omega)$ in terms of the incoming environment particles. To calculate this average, the state of the incoming particle needs to be specified. We first confine the environment particle in a box with periodic boundary conditions. The box volume has a finite value V and the momentum eigenstate in this space is denoted by $|\vec{K}\rangle$. Then we push the box size to the limit of infinity, such that the momentum eigenstate becomes continuous and is denoted by $|\vec{k}\rangle$. Considering the normalization condition, these eigenstates have the following simple well-known connections,

$$|\vec{K}\rangle = \sqrt{\left(\frac{(2\pi)^3}{V}\right)} |\vec{k}\rangle, \quad \frac{(2\pi)^3}{V} \sum = \int d^3\vec{k}. \quad (12)$$

Thus, the state of the incoming particle is described by the density operator,

$$\rho_E = \frac{(2\pi)^3}{V} \sum \mu(\vec{k}) |\vec{K}\rangle \langle \vec{K}|, \quad (13)$$

where the summation runs over the set of momenta that satisfy the periodic boundary condition. $\mu(\vec{k})$ is the wave-number distribution. We assume that the environment is spherically symmetric such that $\mu(\vec{k})$ depends only on the magnitude of \vec{k} . Then the average of the operator $\hat{S}^\dagger(\Omega') \hat{S}(\Omega)$ can be

written as

$$\langle \chi | \hat{S}^\dagger(\Omega') \hat{S}(\Omega) | \chi \rangle \rightarrow \frac{(2\pi)^3}{V} \sum \mu(k) \langle \vec{K} | \hat{S}^\dagger(\Omega') \hat{S}(\Omega) | \vec{K} \rangle. \quad (14)$$

To proceed, the identity $\hat{S} = \hat{I} + i\hat{T}$ is used to express the scattering operator \hat{S} in terms of \hat{T} operator. Recall the definition $\hat{S}(\Omega) = \hat{D}_E(\Omega) \hat{S} \hat{D}_E^\dagger(\Omega)$, the above expression is written as

$$\begin{aligned} \langle \chi(\Omega') | \chi(\Omega) \rangle &\rightarrow 1 - \frac{(2\pi)^3}{V} \int d^3 \vec{k} \mu(k) \langle \vec{k} | \hat{T}^\dagger \hat{T} \\ &- D_E(\Omega') \hat{T}^\dagger D_E^\dagger(\Omega') D_E(\Omega) \hat{T} D_E^\dagger(\Omega) | \vec{k} \rangle, \end{aligned} \quad (15)$$

where the identities $-i\hat{T}^\dagger + i\hat{T} = -\hat{T}^\dagger \hat{T}$, $\int d^3 \vec{k} \mu(k) = 1$, and Eq. (12) are used. While obtaining the above expression, we also used the fact that the environment is spherically symmetric, which is equivalent to state that the environment density operator ρ_E commutes with the environmental rotation $D_E(\Omega)$. For the same reason, the above expression can be written in a more symmetric form,

$$\begin{aligned} \langle \chi(\Omega') | \chi(\Omega) \rangle &\rightarrow 1 - \frac{(2\pi)^3}{2V} \int d^3 \vec{k} \mu(k) \langle \vec{k} | \hat{T}_\Omega^\dagger \hat{T}_\Omega + \hat{T}_\Omega^\dagger \hat{T}_\Omega \\ &- \hat{T}_\Omega^\dagger \hat{T}_\Omega - \hat{T}_\Omega^\dagger \hat{T}_\Omega | \vec{k} \rangle, \end{aligned} \quad (16)$$

where we denote $\hat{T}_\Omega = D_E(\Omega) \hat{T} D_E^\dagger(\Omega)$, which is the rotated \hat{T} operator. Next, we connect the \hat{T} operator with the scattering amplitude by the following familiar formula [21],

$$\langle \vec{k} | \hat{T}_\Omega | \vec{k}' \rangle = -\frac{\hbar^2}{2\pi m} \delta(E - E') f_\Omega(k\hat{k}, k'\hat{k}'), \quad (17)$$

and use the identity operator $\hat{I} = \int d^3 \vec{k}' | \vec{k}' \rangle \langle \vec{k}' |$. After several steps of algebra, we obtain

$$\begin{aligned} \langle \chi(\Omega') | \chi(\Omega) \rangle &\rightarrow 1 - \frac{(2\pi)^3}{2V} \int d^3 \vec{k}' \int d^3 \vec{k} \mu(k) \frac{\hbar^4}{(2\pi m)^2} \delta^2(E - E') \\ &\times \{ f_\Omega^*(\vec{k}', \vec{k}) f_\Omega(\vec{k}', \vec{k}) + f_\Omega^*(\vec{k}', \vec{k}) f_\Omega(\vec{k}, \vec{k}) \\ &- f_\Omega^*(\vec{k}, \vec{k}) f_\Omega(\vec{k}', \vec{k}) - f_\Omega^*(\vec{k}, \vec{k}) f_\Omega(\vec{k}', \vec{k}) \}, \end{aligned} \quad (18)$$

where we are encountered with a squared delta function. Inspired by the usual approach in deriving the Fermi's golden rule, the squared delta function can be evaluated by the following formula [2,15,18],

$$\delta^2(E' - E) = \frac{t}{2\pi \hbar} \delta(E' - E) = \frac{t}{2\pi \hbar} \frac{m}{\hbar^2 k} \delta(k' - k), \quad (19)$$

where the parameter t is interpreted as the time when the interaction is on during the scattering event and is assumed to be much shorter than the system's decoherence time induced by a large number of collisions [2]. Using Eq. (19) and integrating the magnitude of momentum k' , we get

$$\begin{aligned} \langle \chi(\Omega') | \chi(\Omega) \rangle &\rightarrow 1 - \frac{t}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \\ &\times \iint d^2 \hat{k} d^2 \hat{k}' \{ f_\Omega^*(k\hat{k}', k\hat{k}) f_\Omega(k\hat{k}', k\hat{k}) \} \end{aligned}$$

$$\begin{aligned} &+ f_\Omega^*(k\hat{k}', k\hat{k}) f_\Omega(k\hat{k}', k\hat{k}) \\ &- f_\Omega^*(k\hat{k}', k\hat{k}) f_\Omega(k\hat{k}, k\hat{k}) \\ &- f_\Omega^*(k\hat{k}, k\hat{k}) f_\Omega(k\hat{k}', k\hat{k}) \}. \end{aligned} \quad (20)$$

The above expression in the integral can be further simplified. Since the scattering amplitude satisfies $f_\Omega(k\hat{k}', k\hat{k}) = f_\Omega^*(k\hat{k}, k\hat{k}')$ and $k\hat{k}, k\hat{k}'$ are symmetric in swapping the integral index, the double solid angle integral of each term is real. It means $\iint d^2 \hat{k} d^2 \hat{k}' f_\Omega^*(k\hat{k}', k\hat{k}) f_\Omega(k\hat{k}', k\hat{k}) = \iint d^2 \hat{k} d^2 \hat{k}' f_\Omega(k\hat{k}', k\hat{k}) f_\Omega^*(k\hat{k}', k\hat{k})$. Thus the above formula can be expressed in a more symmetric form,

$$\begin{aligned} \langle \chi(\Omega') | \chi(\Omega) \rangle &\rightarrow 1 - \frac{t}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \\ &\times \iint d^2 \hat{k} d^2 \hat{k}' | f_\Omega(k\hat{k}', k\hat{k}) - f_\Omega(k\hat{k}, k\hat{k}') |^2. \end{aligned} \quad (21)$$

Equation (21) gives the suppression of the system density-matrix element by one single elastic scattering event. The result depends on the difference of the elastic scattering amplitudes for different orientations. When $\Omega = \Omega'$, the overlap is trivially one, which indicates no suppression on the diagonal elements of the system density matrix. Next we can proceed to derive the time evolution of the system density matrix. By substituting the above result into Eq. (11) and taking the limit $t \rightarrow 0$, the following formula is obtained,

$$\frac{\partial \rho_S(\Omega, \Omega', t)}{\partial t} = -\Lambda \rho_S(\Omega, \Omega', t), \quad (22)$$

where the factor

$$\begin{aligned} \Lambda &= \frac{1}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \iint d^2 \hat{k} d^2 \hat{k}' | f_\Omega(k\hat{k}', k\hat{k}) \\ &- f_\Omega(k\hat{k}, k\hat{k}') |^2. \end{aligned} \quad (23)$$

The above expression shows an exponential decay in the off-diagonal elements of the system density matrix. Λ is the decay rate. Equation (22) describes the decoherence effect by one environment particle scattering. An ensemble of N particles will build up the decoherence effect in a way that the decoherence rate is multiplied by the number of particles N , thus

$$\begin{aligned} \Lambda &= \frac{N}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \iint d^2 \hat{k} d^2 \hat{k}' | f_\Omega(k\hat{k}', k\hat{k}) \\ &- f_\Omega(k\hat{k}, k\hat{k}') |^2, \end{aligned} \quad (24)$$

where $\frac{\hbar k}{m}$ is the environment particle velocity. Thus, we derive the general expression for the decoherence rate of a quantum rotational system from the elastic scattering model. Equations (22) and (24) are our main results in this section. The expression is general for elastic scattering, since we have not specified any concrete form of the scattering amplitude. Taking into account the spherical symmetry of the environment, we could further rewrite Eq. (24). Denote the scattering amplitude as

$$f_\Omega(k\hat{k}', k\hat{k}) = D_E^\dagger(\Omega) f(k\hat{k}', k\hat{k}) D_E(\Omega). \quad (25)$$

$f(k\hat{k}', k\hat{k})$ is the scattering amplitude for a specific configuration of the system, from which the other scattering amplitude $f_\Omega(k\hat{k}', k\hat{k})$ can be obtained by performing a rotation $D_E(\Omega)$. This greatly simplifies our calculations in the following sections. Several operations yield

$$\Lambda = \frac{N}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \iint d^2 \hat{k} d^2 \hat{k}' |f(k\hat{k}', k\hat{k}) - D_E^\dagger(\omega) f(k\hat{k}', k\hat{k}) D_E(\omega)|^2, \quad (26)$$

where we define $D_E(\omega) = D_E^\dagger(\Omega') D_E(\Omega)$, and ω can be interpreted as the absolute angle distance between the two rotational configurations. Equation (26) shows that the decoherence rate only depends on the absolute angle difference ω of the configurations, which must result for the case for a spherically symmetric environment. Equation (26) could greatly simplify our following evaluations since we can always fix one configuration of the system, and fully use its possible symmetry when choosing the coordinates. The other configuration is obtained by just rotating the absolute angle ω from the fixed configuration.

C. Comparison with translational decoherence

In this section, we briefly compare the results of the preceding section to the decoherence of the translational degrees of freedom. Recall the case of collisional decoherence for a system's translational degrees of freedom [2,14,15], the system density matrix exponentially decays in terms of time

$$\frac{\partial \rho_S(\vec{x}, \vec{x}', t)}{\partial t} \propto -\Lambda \rho_S(\vec{x}, \vec{x}', t). \quad (27)$$

The decoherence factor Λ is given by

$$\Lambda = \frac{N}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \times \iint d^2 \hat{k} d^2 \hat{k}' |f_{\vec{x}}(k\hat{k}', k\hat{k}) - f_{\vec{x}'}(k\hat{k}', k\hat{k})|^2, \quad (28)$$

where $f_{\vec{x}}(k\hat{k}', k\hat{k}) = e^{i\vec{k}\vec{x}} f(k\hat{k}', k\hat{k}) e^{-i\vec{k}\vec{x}'}$. The above formula (28) takes a similar form as Eq. (24), where the decoherence rate of the system density matrix depends on the difference of the scattering amplitudes at different values of the pointer variable. The difference of the scattering amplitudes quantifies the distance measure in translational or orientational space. After some operations, one can show that the above decoherence factor is equivalent to

$$\Lambda = \frac{N}{V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \iint d^2 \hat{k} d^2 \hat{k}' (1 - e^{i\vec{k}(\hat{k}-\hat{k}')(\vec{x}-\vec{x}')}) \times |f(k\hat{k}, k\hat{k}')|^2, \quad (29)$$

which is the familiar form for the translational decoherence rate. More details can be found in Refs. [14,15]. In the long-wavelength limit, one finds that the translational decay rate is proportional to the position difference square. Similarly, we are expecting the rotational decoherence rate to depend on the angular distance in corresponding orientational space, which is shown in the following sections.

III. DECOHERENCE DUE TO SCATTERING OF THERMAL PHOTONS AND MASSIVE PARTICLES

In this section, we will explore the theory of rotational decoherence by calculating the decoherence rate for two different sources of decoherence: thermal photons and massive particles.

A. Thermal photon scattering

We first consider a dielectric ellipsoid immersed in a photon-gas environment. Assuming blackbody radiation at temperature T_E , the average occupation number of photons with energy $\hbar ck$ is given by the Planck distribution, thus the probability distribution of k with N photons in volume V is

$$\mu(k) = \frac{V}{N} \frac{2}{\exp\left(\frac{\hbar ck}{k_B T_E}\right) - 1}, \quad (30)$$

where c is the speed of light. To get the decoherence rate, a key task is to evaluate the scattering amplitude difference. For a dielectric object, the cross section is determined by the scattered radiation from the induced dipole. A detailed discussion can be found in Ref. [22]. If we have an incoming field $\vec{E}_{\text{inc}} = \vec{\xi} E \exp(-ik\hat{k} \cdot \vec{r})$ with a polarization vector $\vec{\xi}$, the far-field approximation gives a scattering amplitude

$$f(k\hat{k}', k\hat{k}) = \frac{k^2}{4\pi \epsilon_0 E} \vec{\xi}' \cdot \vec{p}, \quad (31)$$

where $\vec{\xi}'$ is the polarization of the outgoing radiation, and \vec{p} is the induced dipole moment. The induced dipole moment is given by

$$\vec{p} = \vec{\alpha}_\Omega \cdot \vec{E}_{\text{inc}}, \quad (32)$$

where $\vec{\alpha}_\Omega$ is the polarizability of the ellipsoid with configuration $\Omega = (\alpha, \gamma, \beta)$. According to Eq. (26), we can always choose a configuration with the semiaxis of the ellipsoid aligned with the coordinate axis, such that the polarizability is diagonal

$$\vec{\alpha}_0 = \begin{pmatrix} \alpha_x & 0 & 0 \\ 0 & \alpha_y & 0 \\ 0 & 0 & \alpha_z \end{pmatrix}, \quad (33)$$

where the subscript means the Euler angles are zero for this situation. Then the polarizability with any configuration can be easily derived through the following rotation,

$$\vec{\alpha}_{\Omega'} = R^\dagger(\Omega') \vec{\alpha}_0 R(\Omega'). \quad (34)$$

Now we can calculate the difference of the scattering amplitudes in Eq. (26). Through combining Eqs. (31)–(34), the integral becomes

$$\Lambda = \frac{N}{2V} \int dk k^2 \mu(k) \frac{\hbar k}{m} \times \iint d^2 \hat{k} d^2 \hat{k}' \frac{k^4}{(4\pi \epsilon_0)^2} |\vec{\xi}' \cdot (\vec{\alpha}_0 - \vec{\alpha}_{\Omega'}) \cdot \vec{\xi}|^2, \quad (35)$$

where $\frac{\hbar k}{m} = c$. In order to evaluate the above integral, we should first average over the polarization direction of the incoming and outgoing field. The procedure is simplified by

adopting the following useful identity,

$$\sum_{\lambda} \xi_i^{(\lambda)} \xi_j^{(\lambda)} = \delta_{ij} - \hat{k}_i \cdot \hat{k}_j, \quad (36)$$

where λ is the polarization index and $\{\vec{\xi}^{(1)}, \vec{\xi}^{(2)}, \hat{k}\}$ form an orthogonal basis set. Including the distribution given by Eq. (30), the integral gives the final result

$$\Lambda = 6! \frac{c}{36\epsilon_0^2} \left(\frac{k_B T_E}{\hbar c} \right)^7 \zeta(7) \mathcal{L}, \quad (37)$$

where

$$\zeta(n) = \frac{1}{(n-1)!} \int_0^{\infty} d\chi \frac{\chi^{n-1}}{e^{\chi} - 1}$$

is the Riemann ζ function and $\zeta(7) \simeq 1.00835$, c is the speed of light, and the other parameters are

$$\mathcal{L} = A^2 a_1 + B^2 a_2 + C^2 a_3 + A B a_4 + A C a_5 - B C a_6,$$

$$A = \alpha_x - \alpha_y,$$

$$B = \alpha_x - \alpha_z,$$

$$C = \alpha_y - \alpha_z,$$

$$a_1 = 3 - 3 \cos(2\alpha) \cos(2\gamma) - \cos(2\alpha) \cos(2\beta) \cos(2\gamma) \\ + 4 \cos \beta \sin(2\alpha) \sin(2\gamma),$$

$$a_2 = a_3 = 2 - \cos(2\beta),$$

$$a_4 = a_5 = 2 \cos(2\alpha) \sin^2 \beta + 2 \cos(2\gamma) \sin^2 \beta,$$

$$a_6 = 2 \cos(2\beta). \quad (38)$$

In this expression, the polarizabilities α_x , α_y , and α_z should not be confused with the Euler angle α inside the trigonometric functions. The above equation gives the general expression for the decoherence rate of a dielectric ellipsoid, which depends on the Euler angles and the components of polarizability. This expression gives $\mathcal{L} = 0$ when all angles are zero, as befitting the requirement that decoherence leaves the diagonal elements of the density matrix unchanged.

If the ellipsoid is cylindrically symmetric, the polarizability components $\alpha_x = \alpha_y$. In this case, only one angle dependence is expected in the decoherence rate, and the result emerges from the general expression. Considering a z -axis cylindrical symmetrical ellipsoid, we have $\alpha_x = \alpha_y$. The above result can be reduced to

$$\Lambda = 6! \frac{2c}{9\epsilon_0^2} \left(\frac{k_B T_E}{\hbar c} \right)^7 \zeta(7) (\alpha_x - \alpha_z)^2 \sin^2 \beta, \quad (39)$$

where β is the difference in angle between the two orientations. First, we see that the decoherence rate only depends on the angular difference between the two orientations; the decay rate depends on the sine square of the difference in angles, and it will get its maximal when $\beta = \pi/2$. This is reasonable because, as β increases, the configuration begins to repeat itself when β becomes larger than $\pi/2$. Second, the decoherence rate strongly depends on the temperature. Increasing the temperature will greatly suppress quantum coherence. Also, we find that the temperature dependence for rotational decoherence is two powers lower in T_E than that for center of mass decoherence [2]. Third, if we totally symmetrize the

TABLE I. Estimates of the rotational ($1/\Lambda$) and translational ($1/\mathcal{L}$) decoherence time for different temperatures. The angular separation $\beta = \pi/20$.

Temperature (K)	Rotation (s)	Translation (s)
3	10^{11}	10^{20}
50	10^3	10^9
100	10	10^6
200	10^{-1}	10^3
300	10^{-3}	10^2

system by setting $\alpha_x = \alpha_y = \alpha_z$, the orientational decoherence rate will be equal to zero because the photon scattering cannot distinguish the rotational state of a sphere.

For a specific example, we consider an ellipsoidal nanodiamond, with the size about 100 nm. The nanodiamond shape typically is not elliptical, but the following is an estimate and in the long-wavelength limit the precise shape is not important. In the evaluation, we pick 50 and 75 nm, respectively, as the short and long half axis of the nanodiamond, which gives ellipticity $e = 0.75$. The polarizability satisfies [23]

$$\alpha_i \sim \epsilon_0 V \frac{\epsilon_d - \epsilon_0}{\epsilon_0 + L_i(\epsilon_d - \epsilon_0)}, \quad (40)$$

where $\epsilon_d \sim 6\epsilon_0$ and ϵ_0 are the diamond and vacuum dielectric constant, respectively, and V is the volume of the diamond. L_i , where the index ($i = x, y, z$), is determined by the ellipticity of the nanodiamond. For an ellipsoidal diamond with ellipticity $e = 0.75$, one can get $L_z \sim 0.23$, and $L_x = L_y \sim 0.38$. Thus, the decoherence rate is approximated by $\Lambda \sim 3.2 \times 10^{-14} \left(\frac{T_E}{K} \right)^7 \sin^2 \beta$ (1/s).

Referring to the translational degrees of freedom [20], the decoherence rate is given by

$$\mathcal{L} = 8! \frac{1}{2\pi^3} V^2 c \left(\frac{\epsilon_d - \epsilon_0}{\epsilon_d + 2\epsilon_0} \right)^2 \left(\frac{k_B T_E}{\hbar c} \right)^9 \zeta(9) \Delta x^2, \quad (41)$$

where $\zeta(9) = 1.002$ and V is the volume of the system particle. It is obvious that the ratio of rotational decoherence rate to translational decoherence rate is proportional to $\left(\frac{\hbar c}{k_B T_E r} \right)^2 \sim \frac{1}{(k_{th} r)^2} \gg 1$, where k_{th} is a thermal photon wave number. In order to roughly compare the rotational and translational decoherence rates, we also pick a nanodiamond with a radius $r \sim 50$ nm. The spacial separation is chosen $\Delta x = r \sin \beta$, which is equal to the ‘‘distance’’ the tip of the ellipsoid moves. Thus, the translational decoherence rate is approximated by $\mathcal{L} = 1.5 \times 10^{-23} \left(\frac{T_E}{K} \right)^9 \sin^2 \beta$ (1/s).

Table I shows the rotational ($1/\Lambda$) and translational ($1/\mathcal{L}$) decoherence time scales for several temperatures. First, one can see that the coherence time drops dramatically as the temperature grows from a cold environment to room temperature. Second, for the same temperature, the translational coherence time is much longer than that for rotational degrees of freedom. A simple reason is that, when a nanodiamond is moved a distance x having fixed orientation, the scattering hardly changes because the dielectric polarizability stays the same. However, if we rotate the nanodiamond, the pattern of scattering changes dramatically due to the change of the polarizability. Therefore, it is easier to entangle photons with rotations than with translations.

At last, it is worth mentioning that rotational decoherence is present whenever the different axes have different polarizabilities. A spherical birefringent dielectric will have different polarizabilities α_i in different directions, leading to a nonzero decoherence rate in Eq. (39). Thus, birefringence can be used to control nanoparticles [24,25] but will also lead to rotational decoherence from the asymmetrically scattered photons as discussed in this section.

B. Massive particle scattering

In this section, we will consider massive particles as the source of decoherence in an ellipsoidal system. These types of massive particles typically have a very short de Broglie wavelength, for example, the O_2 molecule at room temperature has a de Broglie wavelength $\lambda_d \sim 10^{-11}$ m. The size of a system, such as a nanodiamond or dust particles, is much larger than λ_d , which trivially indicates that one single scattering could carry away a maximum orientational information. However, as pointed out in Refs. [2,15], it is necessary to determine the lower bound of the decoherence rate. So we still employ the long-wavelength approximation in the following evaluation. The massive particles are assumed to be in thermal equilibrium, which gives the Maxwell-Boltzmann distribution

$$\mu(k) = \left(\frac{\hbar^2}{2\pi m k_B T} \right)^{3/2} \exp\left(-\frac{\hbar^2 k^2}{2m k_B T}\right). \quad (42)$$

Next, we will adopt the Born approximation to evaluate the scattering amplitude. In the Born approximation, the scattering amplitude is given by the following formula,

$$f(\vec{k}', \vec{k}) = -\frac{m}{2\pi \hbar^2} \int d^3 \vec{r} \exp\{-i(\vec{k}' - \vec{k})\vec{r}\} V(\vec{r}), \quad (43)$$

in which $V(\vec{r})$ is the potential of the system. In a real situation, the potential can be very complicated. Here, we consider a cylindrical symmetric ellipsoid which is modeled by the following potential,

$$V(\vec{r}) = D^\dagger(\Omega) V_0(\vec{r}) D(\Omega), \quad (44)$$

where $V_0(\vec{r}) = V_0 \exp\{-a(x^2 + y^2) - bz^2\}$, with its symmetric axis placed at the z direction. The parameters a and b are positive and unequal. Due to the cylindrical symmetry, two Euler angles are enough to specify the orientation. The symmetric axis of the potential $V(\vec{r})$ is in any direction determined by the Euler angles. We first calculate the scattering amplitude Eq. (43) with the above potential $V(\vec{r})$. A convenient way to do the integral is in Cartesian coordinates. We first perform a coordinate rotation $O(x, y, z) \rightarrow \tilde{O}(\tilde{x}, \tilde{y}, \tilde{z})$ to get $V(\vec{r}) \rightarrow \tilde{V}(\vec{r})$, such that the symmetric axis of the potential $\tilde{V}(\vec{r})$ is aligned with the \tilde{z} axis, then calculate the integral in the \tilde{O} coordinate. At last, the final scattering amplitude is obtained by rotating the integral result back to the original coordinate O . Finally, the scattering amplitude is given by

$$f_{\alpha, \beta}(\vec{k}', \vec{k}) = \frac{m V_0 \pi}{2\pi \hbar^2} \frac{\pi}{a} \sqrt{\frac{\pi}{b}} \exp\left\{-\frac{\widetilde{\Delta k_x}^2}{4a} - \frac{\widetilde{\Delta k_y}^2}{4a} - \frac{\widetilde{\Delta k_z}^2}{4b}\right\}, \quad (45)$$

where the vectors $\vec{\widetilde{\Delta k}} = (\widetilde{\Delta k_x}, \widetilde{\Delta k_y}, \widetilde{\Delta k_z})$ and $\vec{\Delta k} = (\Delta k_x, \Delta k_y, \Delta k_z)$ satisfy $\vec{\widetilde{\Delta k}} = R_y^{-1}(\beta) R_z^{-1}(\alpha) \vec{\Delta k}$, and $\Delta k_i =$

$k'_i - k_i$, $\widetilde{\Delta k}_i = \widetilde{k}'_i - \widetilde{k}_i$ ($i = x, y, z$). In the following evaluation, we can Taylor expand Eq. (45) and keep the first-order term in the long-wavelength limit. Substituting Eqs. (42) and (45) into Eq. (26), we get

$$\Lambda = \frac{32\pi}{15\hbar^8} N/V \sqrt{2\pi m^7 (k_b T)^5} \frac{(a-b)^2 V_0^2}{a^4 b^3} \sin^2 \beta. \quad (46)$$

First, we see that the decoherence rate only depends on the polar angle, which specifies the angle difference for the current situation. Second, the parameter $\frac{(a-b)^2 V_0^2}{a^4 b^3}$ is determined by the size and geometry of the system. When we set $a = b$, the system becomes spherically symmetric, which reduces the decoherence rate to zero. Moreover, the rate has a dependence on the two and a half power of the temperature, which is one power higher than that for the case of center of mass decoherence [2]. At last, the rate is also proportional to environment particle density N/V , which is quite reasonable because higher density increases the scattering rate.

IV. CONCLUSION

Decoherence, since the early 1980's, has been used to study a vast array of phenomena ranging from microscopic to cosmological scales [26]. Much effort has been devoted to obtain the pointer states of a given master equation, which is important in establishing the quantum to classical transition. Based on a scattering model [14,15], the decoherence of a mesoscopic system is widely studied for the center of mass motion, where the environmentally distinguished states become exponentially localized wave packets [2,15].

In this paper, we show that the same decoherence effect holds for a mesoscopic quantum system with rotational degrees of freedom. We developed the equations that describe rotational decoherence due to random collisions when a mesoscopic system interacts with an external environment. An environment consisting of a photon gas or massive particles is able to exponentially localize the rotational state, with the decoherence rate proportional to the difference of rotational configurations. Interestingly, the decay rate has a temperature dependence that is different from that for translational decoherence. The rotational decoherence due to photons seems to be faster than translational decoherence by a factor of $1/(k_{th} r)^2 \gg 1$, where k_{th} is a thermal photon wave number and r is the particle size.

The study of rotational decoherence is instrumental for the growing interests in accurate quantum control over a system's internal motion. As more experimental evidence for mesoscopic quantum phenomena is found, considering the decoherence in all degrees of freedom becomes critical. In Ref. [27], the decoherence of center of mass motion induced by the interaction with a system's own internal degrees of freedom is even suggested. For any accurate quantum control or the manufacture of quantum devices, it is extremely important to identify the decoherence mechanism and the corresponding decoherence time. In conclusion, the study of rotational decoherence, together with decoherence with other degrees of freedom, will surely contribute as a useful guidance to future mesoscopic-scale experiments and applications.

ACKNOWLEDGMENTS

The authors thank Prof. K. Hornberger for providing Ref. [17] and Prof. Tongcang Li for helpful discussions. This

work was supported by the National Science Foundation under Grant No. 1404419-PHY.

-
- [1] E. Joos, H. D. Zeh, C. Kiefer, D. J. Giulini, J. Kupsch, and I.-O. Stamatescu, *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer, Berlin, 2013).
- [2] M. Schlosshauer, Decoherence, the measurement problem, and interpretations of quantum mechanics, *Rev. Mod. Phys.* **76**, 1267 (2005).
- [3] W. H. Zurek, Decoherence and the transition from quantum to classical—revisited, [arXiv:quant-ph/0306072](https://arxiv.org/abs/quant-ph/0306072).
- [4] W. H. Zurek, Decoherence, einselection, and the quantum origins of the classical, *Rev. Mod. Phys.* **75**, 715 (2003).
- [5] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2010).
- [6] D. E. Chang, C. Regal, S. Papp, D. Wilson, J. Ye, O. Painter, H. J. Kimble, and P. Zoller, Cavity opto-mechanics using an optically levitated nanosphere, *Proc. Natl. Acad. Sci. USA* **107**, 1005 (2010).
- [7] W. H. Zurek, Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse?, *Phys. Rev. D* **24**, 1516 (1981).
- [8] S. Gerlich, S. Eibenberger, M. Tomandl, S. Nimmrichter, K. Hornberger, P. J. Fagan, J. Tüxen, M. Mayor, and M. Arndt, Quantum interference of large organic molecules, *Nat. Commun.* **2**, 263 (2011).
- [9] A. J. Leggett, S. Chakravarty, A. Dorsey, M. P. Fisher, A. Garg, and W. Zwerger, Dynamics of the dissipative two-state system, *Rev. Mod. Phys.* **59**, 1 (1987).
- [10] Y. Nakamura, Y. A. Pashkin, and J. Tsai, Coherent control of macroscopic quantum states in a single-Cooper-pair box, *Nature (London)* **398**, 786 (1999).
- [11] Y. Nakamura, C. D. Chen, and J. S. Tsai, Spectroscopy of Energy-Level Splitting Between two Macroscopic Quantum States of Charge Coherently Superposed by Josephson Coupling, *Phys. Rev. Lett.* **79**, 2328 (1997).
- [12] O. Romero-Isart, M. L. Juan, R. Quidant, and J. I. Cirac, Toward quantum superposition of living organisms, *New J. Phys.* **12**, 033015 (2010).
- [13] T. Li and Z.-Q. Yin, Quantum superposition, entanglement, and state teleportation of a microorganism on an electromechanical oscillator, *Sci. Bull.* **61**, 163 (2016).
- [14] S. L. Adler, Normalization of collisional decoherence: Squaring the delta function, and an independent cross-check, *J. Phys. A: Math. Gen.* **39**, 14067 (2006).
- [15] K. Hornberger and J. E. Sipe, Collisional decoherence reexamined, *Phys. Rev. A* **68**, 012105 (2003).
- [16] T. M. Hoang, Y. Ma, J. Ahn, J. Bang, F. Robicheaux, Z.-Q. Yin, and T. Li, Torsional Optomechanics of a Levitated Nonspherical Nanoparticle, *Phys. Rev. Lett.* **117**, 123604 (2016).
- [17] T. Fischer, Decoherence of the orientation state, Ph.D. thesis, Duisburg, 2014.
- [18] M. Busse and K. Hornberger, Pointer basis induced by collisional decoherence, *J. Phys. A: Math. Theor.* **43**, 015303 (2009).
- [19] T. Fischer, C. Gneiting, and K. Hornberger, Wigner function for the orientation state, *New J. Phys.* **15**, 063004 (2013).
- [20] E. Joos and H. D. Zeh, The emergence of classical properties through interaction with the environment, *Z. Phys. B* **59**, 223 (1985).
- [21] J. J. Sakurai and J. Napolitano, *Modern Quantum Mechanics* (Addison-Wesley, Boston, 2011).
- [22] J. D. Jackson, *Classical Electrodynamics* (Wiley, New York, 1999).
- [23] C. F. Bohren and D. R. Huffman, *Absorption and Scattering of Light By Small Particles* (Wiley, Hoboken, NJ, 2008).
- [24] Y. Arita, M. Mazilu, and K. Dholakia, Laser-induced rotation and cooling of a trapped microgyroscope in vacuum, *Nat. Commun.* **4**, 2374 (2013).
- [25] A. La Porta and M. D. Wang, Optical Torque Wrench: Angular Trapping, Rotation, and Torque Detection of Quartz Microparticles, *Phys. Rev. Lett.* **92**, 190801 (2004).
- [26] J. B. Hartle and S. W. Hawking, Wave function of the universe, *Phys. Rev. D* **28**, 2960 (1983).
- [27] T. A. Brun and L. Mlodinow, Decoherence by internal degrees of freedom, [arXiv:1512.03845](https://arxiv.org/abs/1512.03845).